

DESCRIPTION

NEW COMPOUND

5 TECHNICAL FIELD

This invention relates to new propanolamine derivatives which is the β_3 adrenergic receptor agonist and salts thereof which are useful as a medicament.

10 DISCLOSURE OF INVENTION

This invention relates to new propanolamine derivatives which is the β_3 adrenergic receptor agonist and salts thereof.

More particularly, it relates to new propanolamine
15 derivatives and salts thereof which have gut selective
sympathomimetic, anti-ulcerous, anti-pancreatitis, lipolytic,
anti-urinary incontinence and anti-pollakiuria activities, to
processes for the preparation thereof, to a pharmaceutical
composition comprising the same and to a method of using the
20 same therapeutically in the treatment and/or prevention of
gastro-intestinal disorders caused by smooth muscle
contractions in human beings or animals, and more
particularly to a method for the treatment and/or prevention
of spasm or hyperanakisia in case of irritable bowel
25 syndrome, gastritis, gastric ulcer, duodenal ulcer,
enteritis, cholecystopathy, cholangitis, urinary calculus and
the like; for the treatment and/or prevention of ulcer such
as gastric ulcer, duodenal ulcer, peptic ulcer, ulcer caused
by non steroidal anti-inflammatory drugs, or the like; for
30 the treatment and/or prevention of dysuria such as
pollakiuria, urinary incontinence or the like in case of
nervous pollakiuria, neurogenic bladder dysfunction,
nocturia, unstable bladder, cystospasm, chronic cystitis,
chronic prostatitis, overflow incontinence, passive
35 incontinence, reflux incontinence, urge incontinence, urinary

stress incontinence or the like; and for the treatment and/or prevention of pancreatitis, obesity, diabetes, glycosuria, hyperlipidemia, hypertension, atherosclerosis, glaucoma, melancholia, depression and the like.


One object of this invention is to provide new and useful propanolamine derivatives and salts thereof which have gut selective sympathomimetic, anti-ulcerous, lipolytic, anti-urinary incontinence and anti-pollakiuria activities.

Another object of this invention is to provide processes for the preparation of said propanolamine derivatives and salts thereof.

A further object of this invention is to provide a pharmaceutical composition comprising, as an active ingredient, said propanolamine derivatives and salts thereof.

Still further object of this invention is to provide a therapeutical method for the treatment and/or prevention of aforesaid diseases in human beings or animals, using said propanolamine derivatives and salts thereof.

The object propanolamine derivatives of this invention are new and can be represented by the following general formula [I] :






wherein


R¹ is aryl which may have one or more suitable substituent(s), heterocyclic group or cyclo(lower)alkyl,

R² is hydrogen or amino protective group,

R³ and R⁴ are independently hydrogen, halogen, hydroxy, amino, nitro, carboxy, protected carboxy, aryl, lower alkyl, hydroxy(lower)alkyl, amino(lower)alkyl, acyloxy(lower)alkyl, acylamino(lower)alkyl, lower alkylamino(lower)alkyl which may have one or more

[illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible]

elimination reaction
of amino protective
group



wherein R^1 , R^2 , R^3 , R^4 , R^5 , A, X and m are each as defined
above, and
 R_a^2 is amino protective group.

In the above and subsequent description of the present
specification, suitable examples of the various definition to
be included within the scope of the invention are explained
in detail in the following.

The term "lower" is intended to mean a group having 1 to
6 carbon atom(s), unless otherwise provided.

The preferable number of the "one or more" in the term
of "one or more suitable substituent(s)" may be 1 to 4.

Suitable example of "halogen" may be fluoro, chloro,
bromo, iodo, and the like.

Suitable example of "lower alkyl" may include straight
or branched one having 1 to 6 carbon atom(s), such as methyl,
ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-
butyl, pentyl, 1-methylpentyl, tert-pentyl, neo-pentyl,
hexyl, isohexyl and the like.

Suitable example of "higher alkyl" may include straight
or branched one having 7 to 20 carbon atoms, such as heptyl,
octyl, 3,5-dimethyloctyl, 3,7-dimethyloctyl, nonyl, decyl,
undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl,

hexadecyl, heptadecyl, octadecyl, nonadecyl, icosyl, and the like.

Suitable "lower alkoxy" and "lower alkoxy" moiety may be a straight or branched one such as methoxy, ethoxy, propoxy, isopropoxy, 1-ethylpropoxy, butoxy, sec-butoxy, tert-butoxy, pentyloxy, neopentyloxy, tert-pentyloxy, hexyloxy, and the like.

Suitable example of "aryl" and "ar" moiety may include phenyl which may have lower alkyl (e.g., phenyl, mesityl, tolyl, etc.), naphthyl, anthryl, and the like, in which the preferred one may be phenyl and naphthyl.

Suitable example of "aroyl" moiety may include benzoyl, toluoyl, naphthoyl, anthrylcarbonyl, and the like, in which the preferred one may be benzoyl and naphthoyl.

Suitable example of "protected carboxy" may be a conventional protecting group such as an esterified carboxy group, or the like, and concrete examples of the ester moiety in said esterified carboxy group may be the ones such as lower alkyl ester [e.g. methyl ester, ethyl ester, propyl ester, isopropyl ester, butyl ester, isobutyl ester, tert-butyl ester, pentyl ester, hexyl ester, 1-cyclopropylethyl ester, etc.] which may have suitable substituent(s), for example, lower alkanoyloxy(lower)alkyl ester [e.g. acetoxymethyl ester, propionyloxymethyl ester, butyryloxymethyl ester, valeryloxymethyl ester, pivaloyloxymethyl ester, 1-acetoxyethyl ester, 1-propionyloxyethyl ester, pivaloyloxyethyl ester, 2-propionyloxyethyl ester, hexanoyloxymethyl ester, etc.], lower alkanesulfonyl(lower)alkyl ester [e.g. 2-mesyloethyl ester, etc.] or mono(or di or tri)halo(lower)alkyl ester [e.g. 2-iodoethyl ester, 2,2,2-trichloroethyl ester, etc.];

higher alkyl ester [e.g. heptyl ester, octyl ester, 3,5-dimethyloctyl ester, 3,7-dimethyloctyl ester, nonyl ester, decyl ester, undecyl ester, dodecyl ester, tridecyl ester, tetradecyl ester, pentadecyl ester, hexadecyl ester, heptadecyl ester, octadecyl ester, nonadecyl ester, adamantyl ester, etc.];

lower alkenyl ester [e.g. (C₂-C₆)alkenyl ester (e.g. vinyl ester, allyl ester, etc.)];

lower alkynyl ester [e.g. (C₂-C₆)alkynyl ester (e.g. ethynyl ester, propynyl ester, etc.)];

ar(lower)alkyl ester which may have one or more suitable substituent(s) [e.g. phenyl(lower)alkyl ester which may have 1 to 4 lower alkoxy, halogen, nitro, hydroxy, lower alkyl, phenyl, or halo(lower)alkyl (e.g. benzyl ester, 4-methoxybenzyl ester, 4-chlorobenzyl ester, 4-nitrobenzyl ester, phenethyl ester, trityl ester, benzhydryl ester, bis(methoxyphenyl)methyl ester, 3,4-dimethoxybenzyl ester, 4-hydroxy-3,5-di-tert-butylbenzyl ester, 4-trifluoromethylbenzyl ester, etc.)];

aryl ester which may have one or more suitable substituent(s) [e.g. phenyl ester which may have 1 to 4 lower alkyl, or halogen (e.g. phenyl ester, 4-chlorophenyl ester, tolyl ester, 4-tert-butylphenyl ester, xylyl ester, mesityl ester, cumenyl ester, etc.)];

cycloalkyloxycarbonyloxy(lower)alkyl ester which may have lower alkyl (e.g., cyclopentyloxycarbonyloxymethyl ester, cyclohexyloxycarbonyloxymethyl ester, cycloheptyloxycarbonyloxymethyl ester, 1-methylcyclohexyloxycarbonyloxymethyl ester, 1-(or 2-)[cyclopentyloxycarbonyloxy]ethyl ester, 1-(or 2-)[cyclohexyloxycarbonyloxy]ethyl ester, 1-(or 2-)-[cycloheptyloxycarbonyloxy]ethyl ester, etc.), etc.);

(5-(lower)alkyl-2-oxo-1,3-dioxol-4-yl)(lower)alkyl ester [e.g., (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl ester,

(5-ethyl-2-oxo-1,3-dioxol-4-yl)methyl ester, (5-propyl-2-oxo-

1,3-dioxol-4-yl)methyl ester, 1-(or 2-)(5-methyl-2-oxo-1,3-dioxol-4-yl)ethyl ester, 1-(or 2-)(5-ethyl-2-oxo-1,3-dioxol-4-yl)ethyl ester, 1-(or 2-)(5-propyl-2-oxo-1,3-dioxol-4-yl)ethyl ester, etc.]; or the like,

in which the preferred one may be lower alkyl ester, lower alkanoyloxy(lower)alkyl ester, ar(lower)alkyl ester which may have one or more suitable substituent(s),

cycloalkyloxycarbonyloxy(lower)alkyl ester which may have lower alkyl, higher alkyl ester, and [5-(lower)alkyl-2-oxo-1,3-dioxol-4-yl](lower)alkyl ester;

and the more preferred one may be methyl ester, ethyl ester, isobutyl ester, butyl ester, pentyl ester, hexyl ester, benzyl ester, 4-trifluoromethylbenzyl ester, 4-chlorobenzyl ester, adamantyl ester, (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl ester, (1-cyclohexyloxycarbonyloxy)ethyl ester and pivaloyloxymethyl ester, and the like, in which the preferred one may be (C₁-C₄)alkyl ester, and the most preferred one may be ethyl ester.

Suitable "acyl group" and "acyl" moiety may include carbamoyl, sulfamoyl, sulfinamoyl, sulfenamoyl, aliphatic acyl group and acyl group containing an aromatic ring, which is referred to as aromatic acyl, or heterocyclic ring, which is referred to as heterocyclic acyl.

Suitable example of said acyl may be illustrated as follows :

carbamoyl; sulfamoyl; sulfinamoyl; sulfenamoyl; aliphatic acyl such as lower or higher alkanoyl (e.g., formyl, acetyl, propanoyl, butanoyl, 2-methylpropanoyl, pentanoyl, 2,2-dimethylpropanoyl, hexanoyl, heptanoyl, octanoyl, nonanoyl, decanoyl, undecanoyl, dodecanoyl, tridecanoyl, tetradecanoyl, pentadecanoyl, hexadecanoyl, heptadecanoyl, octadecanoyl, nonadecanoyl, icosanoyl, etc.);

cyclo(lower)alkylcarbonyl (e.g., cyclopropylcarbonyl,

cyclobutylcarbonyl, cyclopentylcarbonyl, cyclohexylcarbonyl, etc.); protected carboxy such as commonly protected carboxy [e.g., esterified carboxy such as lower or higher alkoxy carbonyl (e.g., methoxy carbonyl, ethoxy carbonyl, propyloxy carbonyl, isopropyloxy carbonyl, t-butoxy carbonyl, t-pentyloxy carbonyl, heptyloxy carbonyl, etc.), etc.], or the like; lower alkyl carbamoyl (e.g., methyl carbamoyl, dimethyl carbamoyl, ethyl carbamoyl, propyl carbamoyl, butyl carbamoyl, pentyl carbamoyl, hexyl carbamoyl, etc.); lower or higher alkyl sulfonyl (e.g., methyl sulfonyl, dimethyl sulfonyl, ethyl sulfonyl, etc.); lower or higher alkoxy sulfonyl (e.g., methoxy sulfonyl, ethoxy sulfonyl, etc.); di-(lower) alkoxy phosphoryl (e.g., dimethoxy phosphoryl, diethoxy phosphoryl, dipropoxy phosphoryl, dibutoxy phosphoryl, dipentyloxy phosphoryl, dihexyloxy phosphoryl, etc.), lower alkyl sulfamoyl (e.g., methyl sulfamoyl, dimethyl sulfamoyl, ethyl sulfamoyl, propyl sulfamoyl, butyl sulfamoyl, pentyl sulfamoyl, hexyl sulfamoyl, etc.), aromatic acyl such as aroyl (e.g., benzoyl, toluoyl, naphthoyl, etc.); ar(lower)alkanoyl [e.g., phenyl(lower)alkanoyl (e.g., phenylacetyl, phenylpropanoyl, phenylbutanoyl, phenylisobutanoyl, phenylpentanoyl, phenylhexanoyl, etc.), naphthyl(lower)alkanoyl (e.g., naphthylacetyl, naphthylpropanoyl, naphthylbutanoyl, etc.), etc.]; ar(lower)alkenoyl [e.g., phenyl(lower)alkenoyl (e.g., phenylpropenoyl, phenylbutenoyl, phenylmethacryloyl, phenylpentanoyl, phenylhexenoyl, etc.), naphthyl(lower)alkenoyl (e.g., naphthylpropenoyl, naphthylbutenoyl, etc.), etc.]; ar(lower)alkoxy carbonyl [e.g., phenyl(lower)alkoxy carbonyl (e.g., benzyloxy carbonyl, etc.), etc.]; aryloxy carbonyl (e.g., phenoxy carbonyl, naphthyloxy carbonyl, etc.); aryloxy(lower)alkanoyl (e.g., phenoxyacetyl,

phenoxypropionyl, etc.);

arylcarbamoyl (e.g., phenylcarbamoyl, etc.);

arylthiocarbamoyl (e.g., phenylthiocarbamoyl, etc.);

arylglyoxyloyl (e.g., phenylglyoxyloyl, naphthylglyoxyloyl,
5 etc.); arylsulfonyl (e.g., phenylsulfonyl, p-tolylsulfonyl,
etc.); or the like.

heterocyclic acyl such as

heterocycliccarbonyl;

10 heterocyclic(lower)alkanoyl (e.g., heterocyclicacetyl,

heterocyclicpropanoyl, heterocyclicbutanoyl,

heterocyclicpentanoyl, heterocyclichexanoyl, etc.);

heterocyclic(lower)alkenoyl (e.g., heterocyclicpropenoyl,

heterocyclicbutenoyl, heterocyclicpentenoyl,

15 heterocyclichexanoyl, etc.); heterocyclicglyoxyloyl;

or the like; and the like.

Suitable example of "heterocyclic group" and
"heterocyclic" moiety may include

20 unsaturated 3 to 8-membered (more preferably 5 or 6-
membered) heteromonocyclic group containing 1 to 4 nitrogen
atom(s), for example, pyrrolyl, pyrrolinyl, imidazolyl,
pyrazolyl, pyridyl, dihydropyridyl, pyrimidyl, pyrazinyl,
pyridazinyl, triazolyl (e.g., 4H-1,2,4-triazolyl, 1H-1,2,3-
triazolyl, 2H-1,2,3-triazolyl, etc.), tetrazolyl (e.g., 1H-
25 tetrazolyl, 2H-tetrazolyl, etc.), etc.;

saturated 3 to 8-membered (more preferably 5 or 6-
membered) heteromonocyclic group containing 1 to 4 nitrogen
atom(s), for example, pyrrolidinyl, imidazolidinyl,
piperidyl, piperazinyl, etc.;

30 unsaturated condensed heterocyclic group containing 1 to
4 nitrogen atom(s), for example, indolyl, dihydroindolyl,
isoindolyl, indolinyl, indolizinyll, benzimidazolyl, quinolyl,
isoquinolyl, indazolyl, benzotriazolyl, etc.;

35 unsaturated 3 to 8-membered (more preferably 5 or 6-
membered) heteromonocyclic group containing 1 to 2 oxygen

atom(s) and 1 to 3 nitrogen atom(s), for example, oxazolyl, isoxazolyl, oxadiazolyl (e.g., 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl, etc.), etc.;

5 saturated 3 to 8-membered (more preferably 5 or 6-membered) heteromonocyclic group containing 1 to 2 oxygen atom(s) and 1 to 3 nitrogen atom(s), for example, morpholino, sydnonyl, etc.;

10 unsaturated condensed heterocyclic group containing 1 to 2 oxygen atom(s) and 1 to 3 nitrogen atom(s), for example, benzoxazolyl, benzoxadiazolyl, etc.;

15 unsaturated 3 to 8-membered (more preferably 5 or 6-membered) heteromonocyclic group containing 1 to 2 sulfur atom(s) and 1 to 3 nitrogen atom(s), for example, thiazolyl, isothiazolyl, thiadiazolyl (e.g., 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, etc.), dihydrothiazinyl, etc.;

20 saturated 3 to 8-membered (more preferably 5 or 6-membered) heteromonocyclic group containing 1 to 2 sulfur atom(s) and 1 to 3 nitrogen atom(s), for example, thiazolidinyl, etc.;

25 unsaturated 3 to 8-membered (more preferably 5 or 6-membered) heteromonocyclic group containing 1 to 2 sulfur atom(s), for example, thienyl, dihydrodithiinyl, dihydrodithionyl, etc.;

30 unsaturated condensed heterocyclic group containing 1 to 2 sulfur atom(s) and 1 to 3 nitrogen atom(s), for example, benzothiazolyl, benzothiadiazolyl, imidazothiadiazolyl, etc.;

35 unsaturated 3 to 8-membered (more preferably 5 or 6-membered) heteromonocyclic group containing an oxygen atom, for example, furyl, etc.;

saturated 3 to 8-membered (more preferably 5 or 6-membered) heteromonocyclic group containing an oxygen atom, for example, tetrahydrofuran, tetrahydropyran, etc.;

unsaturated 3 to 8-membered (more preferably 5 or 6-membered) heteromonocyclic group containing an oxygen atom

and 1 to 2 sulfur atom(s), for example, dihydrooxathiinyl, etc.;

unsaturated condensed heterocyclic group containing 1 to 2 sulfur atom(s), for example, benzothienyl, benzodithiinyl, etc.;

unsaturated condensed heterocyclic group containing an oxygen atom and 1 to 2 sulfur atom(s), for example, benzoxathiinyl, etc.; and the like, and the abovementioned "heterocyclic group" and "heterocyclic" moiety may have one or more suitable substituent(s) such as amino, oxo, cyano, aryl, ar(lower)alkyl, heterocyclic group.

Suitable example of "cyclo(lower)alkyl" may include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and the like, in which the preferred one may be cyclo(C₄-C₆)alkyl, and the most preferred one may be cyclopentyl and cyclohexyl.

Suitable example of "amino protective group" moiety may be common amino protective group such as acyl, for example, substituted or unsubstituted lower alkanoyl [e.g. formyl, acetyl, propionyl, trifluoroacetyl, etc.], phthaloyl, lower alkoxy carbonyl [e.g. tert-butoxycarbonyl, tert-amylloxycarbonyl, etc.], substituted or unsubstituted aralkyloxycarbonyl [e.g. benzyloxycarbonyl, p-nitrobenzyloxycarbonyl, etc.], substituted or unsubstituted arenesulfonyl [e.g. benzenesulfonyl, tosyl, etc.], nitrophenylsulfenyl, ar(lower)alkyl [e.g. trityl, benzyl, etc.], and the like, in which preferable one may be lower alkoxy carbonyl and phenyl(lower)alkyl, and the most preferred one may be tert-butoxycarbonyl and benzyl.

Suitable example of "aryl" moiety in the term of "aryl which may have one or more suitable substituent(s)" in R¹ can be referred to aforementioned "aryl", in which the preferred one may be phenyl.

Suitable example of "suitable substituent(s)" moiety in the term of "aryl which may have one or more suitable substituent(s)" may include hydroxy, halogen, lower alkylsulfonylamino, lower alkanoylamino, and the like.

5 Suitable example of "lower alkylsulfonylamino" may include methylsulfonylamino, ethylsulfonylamino, propylsulfonylamino, butylsulfonylamino, pentylsulfonylamino, hexylsulfonylamino, and the like, in which the preferred one may be (C₁-C₄)alkylsulfonylamino, and the most preferred one
10 may be methylsulfonylamino.

 Suitable example of "lower alkanoylamino" may include formylamino, acetylamino, propanoylamino, butanoylamino, 2-methylpropanoylamino, pentanoylamino, hexanoylamino, and the like, in which the preferred one may be (C₁-C₄)-
15 alkanoylamino, and the most preferred one may be acetylamino.

 Suitable example of "heterocyclic group" in R¹ can be referred to aforementioned "heterocyclic group", in which the preferred one may be unsaturated 3 to 8-membered
20 heteromonocyclic group containing 1 to 4 nitrogen atom(s) which may have one or more suitable substituent(s), and unsaturated condensed heterocyclic group containing 1 to 4 nitrogen atom(s) which may have one or more suitable substituent(s), and the most preferred one may be pyridyl,
25 aminopyridyl, indolyl and benzimidazolone.

 Suitable example of "cyclo(lower)alkyl" in R¹ can be referred to aforementioned "cyclo(lower)alkyl", in which the preferred one may be cyclo(C₄-C₆)alkyl, and the most
30 preferred one may be cyclopentyl.

 Suitable example of "mono or di lower alkylamino" may include methylamino, dimethylamino, ethylamino, diethylamino, propylamino, isopropylamino, butylamino, methylbutylamino, sec-butylamino, 1-methylpentylamino, ethylpentylamino,
35 hexylamino, and the like, in which the preferred one may be

mono or di(C₁-C₄)alkylamino, and the most preferred one may be dimethylamino.

Suitable example of "hydroxy(lower)alkyl" may include hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, hydroxypentyl, hydroxyhexyl, and the like, in which the preferred one may be hydroxy(C₁-C₄)alkyl, and the most preferred one may be hydroxymethyl.

Suitable example of "amino(lower)alkyl" may include aminomethyl, aminoethyl, aminopropyl, aminobutyl, aminopentyl, aminohexyl, and the like, in which the preferred one may be amino(C₁-C₄)alkyl, and the most preferred one may be aminomethyl.

Suitable example of "acyl" moiety in the term of "acyloxy(lower)alkyl" can be referred to aforementioned "acyl", in which the preferred one may be lower alkanoyl, and the most preferred one may be acetyl.

Suitable example of "acyloxy(lower)alkyl" may include lower alkanoyloxy(lower)alkyl, in which the preferred one may be (C₁-C₄)alkanoyloxy(C₁-C₄)alkyl, and the most preferred one may be acetyloxymethyl.

Suitable example of "acyl" moiety in the term of "acylamino(lower)alkyl" can be referred to aforementioned "acyl", in which the preferred one may be lower alkanoyl, aroyl, carbamoyl, lower alkylcarbamoyl, lower alkylsulfonyl, and arylsulfonyl.

Suitable example of "acylamino(lower)alkyl" may include lower alkanoylamino(lower)alkyl, aroylamino(lower)alkyl, carbamoylamino(lower)alkyl, lower alkylcarbamoylamino-(lower)alkyl, lower alkylsulfonylamino(lower)alkyl, arylsulfonylamino(lower)alkyl and the like, in which the preferred one may be (C₁-C₄)alkanoylamino(C₁-C₄)alkyl,

benzoylamino(C₁-C₄)alkyl, carbamoylamino(C₁-C₄)alkyl,
 (C₁-C₄)alkylsulfonylamino(C₁-C₄)alkyl and
 phenylsulfonylamino(C₁-C₄)alkyl, and the most referred one
 may be acetylaminomethyl, benzoylaminomethyl,
 5 carbamoylaminomethyl, butylcarbamoylaminomethyl,
 methylsulfonylaminomethyl and phenylsulfonylaminomethyl.

Suitable example of "suitable substituent" moiety in the
 term of "lower alkylamino(lower)alkyl which may have one or
 10 more suitable substituent(s)" may include lower alkyl and
 carboxy, in which the preferred one may be methyl and
 carboxy.

Suitable example of "lower alkylamino(lower)alkyl which
 may have one or more suitable substituent(s)" may include
 15 di(lower)alkylamino(lower)alkyl, carboxy(lower)alkylamino-
 (lower)alkyl, and the like, in which the preferred one may be
 di(C₁-C₄)alkylamino(C₁-C₄)alkyl and carboxy(C₁-C₄)-
 alkylamino(C₁-C₄)alkyl, and the most preferred one may be
 dimethylaminomethyl and carboxyethylaminomethyl.

Suitable example of "acyl" moiety in the term of
 "acylamino" can be referred to aforementioned "acyl" moiety,
 in which the preferred one may be lower alkanoyl, aroyl,
 lower alkoxycarbonyl, aryloxycarbonyl, lower alkylsulfonyl,
 phenylsulfonyl, sulfamoyl, lower alkylsulfamoyl, and the most
 25 preferred one may be acetyl, benzoyl, methoxycarbonyl,
 phenoxycarbonyl, methylsulfonyl, phenylsulfonyl, sulfamoyl
 and dimethylsulfamoyl.

Suitable example of "acylamino" may be acetylamino,
 benzoylamino, methoxycarbonylamino, phenoxycarbonylamino,
 30 methylsulfonylamino, phenylsulfonylamino, sulfamoylamino and
 dimethylsulfamoylamino.

Suitable example of "acyl group" can be referred to
 aforementioned "acyl group", in which the preferred one may
 35 be carbamoyl, lower alkylcarbamoyl, arylcarbamoyl, and the

most preferred one may be carbamoy, methylcarbamoyl, dimethylcarbamoyl and phenylcarbamoyl.

Suitable example of "lower alkoxy" can be referred to
5 aforementioned "lower alkoxy", in which the preferred one may be methoxy, ethoxy, propoxy, isopropoxy and benzyloxy.

Suitable example of "halo(lower)alkoxy" may be
10 fluoro(lower)alkoxy, chloro(lower)alkoxy, bromo(lower)alkoxy, iodo(lower)alkoxy, and the like, in which the preferred one may be fluoro(C₁-C₄)alkoxy, chloro(C₁-C₄)alkoxy, bromo(C₁-C₄)alkoxy, iodo(C₁-C₄)alkoxy, and the most preferred one may be fluoromethoxy.

15 Suitable example of "lower alkenyloxy" may include vinyloxy, 1-(or 2-)propenyloxy, 1-(or 2- or 3-)butenyloxy, 1-(or 2- or 3- or 4-)pentenyloxy, 1-(or 2- or 3- or 4- or 5-)hexenyloxy, and the like, in which the preferred one may be (C₂-C₄)alkenyloxy, and the most preferred one may be 2-propenyloxy.
20

Suitable example of "lower alkoxy(lower)alkoxy" may include methoxymethoxy, methoxyethoxy, methoxypropoxy, ethoxypropoxy, ethoxyethoxy, propoxymethoxy, butoxymethoxy,
25 pentyloxymethoxy, hexyloxymethoxy, hexyloxyethoxy, and the like, in which the preferred one may be (C₁-C₄)alkoxy-(C₁-C₄)alkoxy, and the most preferred one may be methoxyethoxy.

30 Suitable example of "aryloxy" may include phenoxy, mesityloxy, tolyloxy, naphthyloxy, anthryloxy, and the like, in which the preferred one may be phenoxy.

Suitable example of "cyclo(lower)alkyloxy" may include
35 cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy,

and the like, in which the preferred one may be cyclopentyloxy.

Suitable example of "heterocyclic" moiety in the term of "heterocyclicoxy" can be referred to aforementioned "heterocyclic" moiety, in which the preferred one may be unsaturated 3 to 8-membered heteromonocyclic group containing 1 to 4 nitrogen atoms(s) which may have a suitable substituent, and the most preferred one may be pyridyl having cyano.

Suitable example of "heterocyclicoxy" may be pyridyloxy having cyano.

Suitable example of "ar(lower)alkyloxy" may include phenyl(lower)alkyloxy, mesityl(lower)alkyloxy, tolyl(lower)alkyloxy, naphthyl(lower)alkyloxy, anthryl(lower)alkyloxy, and the like, in which the preferred one may be phenyl(C₁-C₄)alkyloxy, and the most preferred one may be benzyloxy and phenetyloxy.

Suitable example of "acyl" moiety in the term of "acyloxy" can be referred to aforementioned "acyl" moiety, in which the preferred one may be carbamoyl and mono or di-lower alkylsulfamoyl, and the most preferred one may be carbamoyl and dimethylsulfamoyl.

Suitable example of "acyloxy" may be carbamoyloxy and mono or di lower alkylsulfamoyloxy and the most preferred one may be carbamoyloxy and dimethylsulfamoyloxy.

Suitable example of "lower alkoxy" moiety in the term of "acyl(lower)alkoxy" can be referred to aforementioned "lower alkoxy", in which the preferred one may be (C₁-C₄)alkoxy, and the most preferred one may be methoxy.

Suitable example of "acyl" moiety in the term of "acyl(lower)alkoxy" may include the aforementioned "acyl"

moiety; lower alkylcarbamoyle which may have one or more suitable substituent(s) selected from the group consisting of hydroxy, lower alkoxy, lower alkyl, ar(lower)alkyl, protected carboxy, carboxy, mono or di lower alkylamino, lower alkylthio, halo(lower)alkyl, aryl which may have one or more suitable substituent(s), and heterocyclic group; arylcarbamoyle which may have one or more suitable substituent(s) selected from the group consisting of lower alkyl, higher alkyl, halogen, halo(lower)alkyl, mono or di-lower alkylamino, lower alkoxy, halo(lower)alkoxy and nitro; heterocycliccarbamoyle which may have one or more suitable substituent(s) selected from the group consisting of ar(lower)alkyl, aryl and heterocyclic(lower)alkyl which may be substituted with heterocyclic group; heterocycliccarbonyl which may have one or more suitable substituent(s) selected from the group consisting of ar(lower)alkyl and heterocyclic group which may be substituted with one or more suitable substituent(s); cyclo(lower)alkylcarbamoyle; guanidinocarbonyl; and the like.

Suitable example of "lower alkylcarbamoyle" may include methylcarbamoyle, ethylcarbamoyle, propylcarbamoyle, isopropylcarbamoyle, butylcarbamoyle, 1-methylbutylcarbamoyle, isobutylcarbamoyle, sec-butylcarbamoyle, tert-butylcarbamoyle, pentylcarbamoyle, isopentylcarbamoyle, tert-pentylcarbamoyle, 1-methylpentylcarbamoyle, neopentylcarbamoyle, hexylcarbamoyle, isohexylcarbamoyle, and the like, in which the preferred one may be methylcarbamoyle, ethylcarbamoyle, propylcarbamoyle, butylcarbamoyle, sec-butylcarbamoyle, isobutylcarbamoyle, 1-methylbutylcarbamoyle, and hexylcarbamoyle.

Suitable example of "lower alkoxy" in the suitable substituent(s) of the term "lower alkylcarbamoyle which may have one or more suitable substituent(s)" can be referred to aforementioned "lower alkoxy", in which the preferred one may be (C₁-C₄)alkoxy, and the most preferred one may be methoxy,

ethoxy and propoxy.

Suitable example of "lower alkyl" in the suitable substituent(s) of the term "lower alkylcarbamoyl which may have one or more suitable substituent(s)" can be referred to aforementioned "lower alkyl", in which the preferred one may be (C₁-C₄)alkyl, and the most preferred one may be methyl and ethyl.

Suitable example of "ar(lower)alkyl" in the suitable substituent(s) of the term "lower alkylcarbamoyl which may have one or more suitable substituent(s)" may include "mono- or di- or tri-phenyl(lower)alkyl" such as benzyl, phenethyl, phenylpropyl, phenylbutyl, phenylpentyl, phenylhexyl, benzhydryl, trityl, and the like, in which the preferred one may be benzyl.

Suitable example of "protected carboxy" in the suitable substituent(s) of the term "lower alkylcarbamoyl which may have one or more suitable substituent(s)" can be referred to aforementioned "protected carboxy", in which the preferred one may be (C₁-C₄)alkyl ester, and the most preferred one may be ethyl ester.

Suitable example of "mono or di-lower alkylamino" in the suitable substituent(s) of the term "lower alkylcarbamoyl which may have one or more suitable substituent(s)" may include methylamino, dimethylamino, ethylamino, diethylamino, propylamino, isopropylamino, butylamino, isobutylamino, sec-butylamino, tert-butylamino, pentylamino, isopentylamino, tert-pentylamino, 1-methylpentylamino, neopentylamino, hexylamino, isohexylamino, and the like, in which the preferred one may be di(C₁-C₄)alkylamino, and the most preferred one may be dimethylamino and diethylamino.

Suitable example of "lower alkylthio" in the suitable substituent(s) of the term "lower alkylcarbamoyl which may have one or more suitable substituent(s)" may include methylthio, ethylthio, propylthio, butylthio, pentylthio, and

the like, in which the preferred one may be (C₁-C₄)alkylthio, and the most preferred one may be methylthio.

Suitable example of "halo(lower)alkyl" in the suitable substituent(s) of the term "lower alkylcarbamoyl which may have one or more suitable substituent(s)" may include tri-halo(lower)alkyl such as trichloromethyl, trichloroethyl, trichloropropyl, trifluoromethyl, trifluoroethyl, trifluoropropyl, tribromomethyl, tribromoethyl, and the like, in which the preferred one may be tri-halo(C₁-C₄)alkyl, and the most preferred one may be trifluoromethyl and trifluoroethyl.

Suitable example of "aryl" in the term of "aryl which may have one or more suitable substituent(s)" can be referred to aforementioned "aryl", in which the preferred one may be phenyl.

Suitable example of "suitable substituent(s)" in the term of "aryl which may have one or more suitable substituent(s)" may include mono or di(lower)alkylamino-(lower)alkyl (e.g., methylaminomethyl, dimethylaminomethyl, dimethylaminoethyl, diethylaminoethyl, dimethylaminopropyl, propylaminopropyl, butylaminoethyl, diethylaminopentyl, etc.), in which the preferred one may be mono or di(C₁-C₄)alkylamino(C₁-C₄)alkyl, and the most preferred one may be dimethylaminomethyl.

Suitable example of "heterocyclic group" can be referred to aforementioned "heterocyclic group", in which the preferred one may be saturated or unsaturated 3 to 8-membered heteromonocyclic group containing 1 to 4 nitrogen atom(s) and saturated 3 to 8-membered heteromonocyclic group containing 1 to 2 oxygen atom(s) and 1 to 3 nitrogen atom(s), and the most preferred one may be pyridyl, imidazolyl and morpholino.

Suitable example of "aryl" moiety in the term of "arylcarbamoyl which may have one or more suitable substituent(s)" can be referred to aforementioned "aryl", in

which the preferred one may be phenyl.

Suitable example of "lower alkyl" in the suitable substituent(s) of the term "arylcarbamoyl which may have one or more suitable substituent(s)" can be referred to
5 aforementioned "lower alkyl", in which the preferred one may be (C₁-C₄)alkyl, and the most preferred one may be methyl and butyl.

Suitable example of "higher alkyl" in the suitable substituent(s) of the term "arylcarbamoyl which may have one
10 or more suitable substituent(s)" can be referred to aforementioned "higher alkyl", in which the preferred one may be (C₇-C₁₀)alkyl, and the most preferred one may be octyl.

Suitable example of "halo(lower)alkyl" in the suitable substituent(s) of the term "arylcarbamoyl which may have one
15 or more suitable substituent(s)" may include tri-halo(lower)alkyl such as trichloromethyl, trichloroethyl, trichloropropyl, trifluoromethyl, trifluoroethyl, trifluoropropyl, tribromomethyl, tribromoethyl, and the like, in which the preferred one may be tri-halo(C₁-C₄)alkyl, and
20 the most preferred one may be trifluoromethyl.

Suitable example of "mono or di-lower alkylamino" in the suitable substituent(s) of the term "arylcarbamoyl which may have one or more suitable substituent(s)" may include
25 methylamino, dimethylamino, ethylamino, diethylamino, propylamino, isopropylamino, butylamino, isobutylamino, sec-butylamino, tert-butylamino, pentylamino, isopentylamino, tert-pentylamino, 1-methylpentylamino, neopentylamino, hexylamino, isohexylamino, and the like, in which the
30 preferred one may be di(C₁-C₄)alkylamino, and the most preferred one may be dimethylamino.

Suitable example of "lower alkoxy" in the suitable substituent(s) of the term "arylcarbamoyl which may have one or more suitable substituent(s)" can be referred to
35 aforementioned "lower alkoxy", in which the preferred one may

be (C₁-C₄)alkoxy, and the most preferred one may be methoxy.

Suitable example of "halo(lower)alkoxy" in the suitable substituent(s) of the term "arylcarbamoyl which may have one or more suitable substituent(s)" may include fluoromethoxy, fluoroethoxy, trifluoromethoxy, chloromethoxy, chloroethoxy, trichloroethoxy, bromomethoxy, iodoethoxy, and the like, in which the preferred one may be halo(C₁-C₄)alkoxy, and the most preferred one may be fluoromethoxy.

Suitable example of "heterocyclic" moiety in the term of "heterocycliccarbamoyl which may have one or more suitable substituent(s)" can be referred to aforementioned "heterocyclic" moiety, in which the preferred one may be saturated or unsaturated 3 to 8-membered heteromonocyclic group containing 1 to 4 nitrogen atom(s), unsaturated condensed heterocyclic group containing 1 to 4 nitrogen atom(s) and unsaturated 3 to 8-membered heteromonocyclic group containing 1 to 2 sulfur atom(s) and 1 to 3 nitrogen atom(s), and the most preferred one may be piperidyl, tetrazolyl, indolyl and thiazolyl.

Suitable example of "ar(lower)alkyl" in the suitable substituent(s) of the term "heterocycliccarbamoyl which may have one or more suitable substituent(s)" may include "mono- or di- or tri-phenyl(lower)alkyl" such as benzyl, phenethyl, phenylpropyl, phenylbutyl, phenylpentyl, phenylhexyl, benzhydryl, trityl, and the like, in which the preferred one may be benzyl.

Suitable example of "lower alkyl" moiety in the term of "heterocyclic(lower)alkyl" in the suitable substituent(s) of the term "heterocycliccarbamoyl which may have one or more suitable substituent(s)" can be referred to aforementioned "lower alkyl", in which the preferred one may be (C₁-C₄)alkyl, and the most preferred one may be methyl.

Suitable example of "heterocyclic" moiety in the term of "heterocyclic(lower)alkyl" in the suitable substituent(s) of

the term "heterocycliccarbamoyl which may have one or more suitable substituent(s)" can be referred to aforementioned "heterocyclic", in which the preferred one may be saturated 3 to 8-membered heteromonocyclic group containing 1 to 4 nitrogen atom(s), and the most preferred one may be piperidyl.

Suitable example of "heterocyclic group" in the term of "heterocyclic(lower)alkyl which may be substituted with heterocyclic group" in the suitable substituent(s) of the term "heterocycliccarbamoyl which may have one or more suitable substituent(s)" can be referred to aforementioned "heterocyclic group", in which the preferred one may be unsaturated condensed heterocyclic group containing 1 to 4 nitrogen atom(s), and the most preferred one may be indolyl.

Suitable example of "heterocyclic" moiety in the term of "heterocycliccarbonyl which may have one or more suitable substituent(s)" can be referred to aforementioned "heterocyclic" moiety, in which the preferred one may be saturated 3 to 8-membered heteromonocyclic group containing 1 to 4 nitrogen atom(s) and unsaturated condensed heterocyclic group containing 1 to 4 nitrogen atom(s), and the most preferred one may be piperidyl, piperazinyl and dihydroindolyl.

Suitable example of "ar(lower)alkyl" in the suitable substituent(s) of the term "heterocycliccarbonyl which may have one or more suitable substituent(s)" may include "mono- or di- or tri-phenyl(lower)alkyl" such as benzyl, phenethyl, phenylpropyl, phenylbutyl, phenylpentyl, phenylhexyl, benzhydryl, trityl, and the like, in which the preferred one may be benzhydryl.

Suitable example of "heterocyclic group" moiety in the term "heterocyclic group which may have one or more suitable substituent(s)" in the suitable substituent(s) of the term "heterocycliccarbonyl which may have one or more suitable

substituent(s)" can be referred to aforementioned
 "heterocyclic group", in which the preferred one may be
 unsaturated 3 to 8-membered heteromonocyclic group containing
 1 to 4 nitrogen atom(s), and the most preferred one may be
 5 pyridazinyl.

Suitable example of "suitable substituent(s)" in the
 term of "heterocyclic group which may be substituted with one
 or more suitable substituent(s)" may include oxo and aryl, in
 which the preferred one may be oxo and phenyl.

Suitable example of "cyclo(lower)alkylcarbamoyle" may
 include cyclopropylcarbamoyle, cyclobutylcarbamoyle,
 cyclopentylcarbamoyle, cyclohexylcarbamoyle, and the like, in
 which the preferred one may be cyclo(C₄-C₆)alkylcarbamoyle,
 15 and the most preferred one may be cyclohexylcarbamoyle.

Suitable example of "lower alkyl" in R⁵ can be referred
 to aforementioned "lower alkyl", in which the preferred one
 may be (C₁-C₄)alkyl, and the most preferred one may be
 20 methyl.

Suitable example of "acyl(lower)alkoxy" may include
 carbamoyle(lower)alkoxy, lower alkylcarbamoyle(lower)alkoxy,
 cyclo(lower)alkylcarbamoyle(lower)alkoxy,
 25 arylcarbamoyle(lower)alkoxy, heterocycliccarbamoyle(lower)-
 alkoxy, N-lower alkyl-lower alkylcarbamoyle(lower)alkoxy,
 ar(lower)alkylcarbamoyle(lower)alkoxy, lower
 alkoxy(lower)alkylcarbamoyle(lower)alkoxy, lower
 alkylthio(lower)alkylcarbamoyle(lower)alkoxy,
 30 di(lower)alkylamino(lower)alkylcarbamoyle(lower)alkoxy,
 heterocycliccarbonyl(lower)alkoxy,
 guanidinocarbonyl(lower)alkoxy,
 hydroxy(lower)alkylcarbamoyle(lower)alkoxy,
 halo(lower)alkylcarbamoyle(lower)alkoxy, protected
 35 carboxy(lower)alkylcarbamoyle(lower)alkoxy having protected

carboxy, heterocyclic(lower)alkylcarbamoyl(lower)alkoxy,
heterocycliccarbamoyl(lower)alkoxy having aryl(lower)alkyl,
arylcarbamoyl(lower)alkoxy having higher alkyl,
arylcarbamoyl(lower)alkoxy having lower alkyl,
5 arylcarbamoyl(lower)alkoxy having halo(lower)alkyl,
arylcarbamoyl(lower)alkoxy having nitro,
arylcarbamoyl(lower)alkoxy having halogen,
arylcarbamoyl(lower)alkoxy having
di(lower)alkylamino(lower)alkyl,
10 heterocycliccarbonyl(lower)alkoxy having benzhydryl, lower
alkylcarbamoyl(lower)alkoxy having benzhydryl,
heterocycliccarbamoyl(lower)alkoxy having
heterocyclic(lower)alkyl substituted with heterocyclic group,
heterocycliccarbonyl(lower)alkoxy having heterocyclic group
15 substituted with oxo and aryl,
N-lower alkyl-ar(lower)alkylcarbamoyl(lower)alkoxy,
N-ar(lower)alkyl-ar(lower)alkylcarbamoyl(lower)alkoxy,
N-lower alkyl-heterocyclic(lower)alkylcarbamoyl(lower)alkoxy,
arylcarbamoyl(lower)alkoxy having lower alkyl,
20 arylcarbamoyl(lower)alkoxy having lower alkoxy,
arylcarbamoyl(lower)alkoxy having halogen,
arylcarbamoyl(lower)alkoxy having halo(lower)alkoxy,
arylcarbamoyl(lower)alkoxy having di(lower)alkylamino,
N-aryl-aryl(lower)alkylcarbamoyl(lower)alkoxy, N-lower alkyl-
25 cyclo(lower)alkylcarbamoyl(lower)alkoxy,

in which the preferred one may be carbamoyl(C₁-C₄)-
alkoxy, (C₁-C₆)alkylcarbamoyl(C₁-C₄)alkoxy, cyclo(C₄-C₆)-
alkylcarbamoyl(C₁-C₄)alkoxy, phenylcarbamoyl(C₁-C₄)alkoxy,
carbamoyl(C₁-C₄)alkoxy substituted with unsaturated 3 to 8-
30 membered heteromonocyclic group containing 1 to 4 nitrogen
atom(s), carbamoyl(C₁-C₄)alkoxy substituted with unsaturated
condensed heterocyclic group containing 1 to 4 nitrogen
atom(s), N-(C₁-C₄)alkyl-(C₁-C₆)alkylcarbamoyl(C₁-C₄)alkoxy,
phenyl(C₁-C₄)alkylcarbamoyl(C₁-C₄)alkoxy,
35 (C₁-C₄)alkoxy(C₁-C₄)alkylcarbamoyl(C₁-C₄)alkoxy,

(C₁-C₄)alkylthio(C₁-C₄)alkylcarbamoyl(C₁-C₄)alkoxy,
 di(C₁-C₄)alkylamino(C₁-C₄)alkylcarbamoyl(C₁-C₄)alkoxy,
 carbonyl(lower)alkoxy substituted with saturated 3 to 8-
 membered heteromonocyclic group containing 1 to 4 nitrogen
 5 atom(s), guanidinocarbamoyl(C₁-C₄)alkoxy,
 hydroxy(C₁-C₄)alkylcarbamoyl(C₁-C₄)alkoxy,
 tri-halo(C₁-C₄)alkylcarbamoyl(C₁-C₄)alkoxy, esterified
 carboxy(C₁-C₄)alkylcarbamoyl(C₁-C₄)alkoxy having esterified
 carboxy, (C₁-C₄)alkylcarbamoyl(C₁-C₄)alkoxy substituted with
 10 saturated 3 to 8-membered heteromonocyclic group containing 1
 to 2 oxygen atom(s) and 1 to 3 nitrogen atom(s),
 (C₁-C₄)alkylcarbamoyl(C₁-C₄)alkoxy substituted with
 unsaturated 3 to 8-membered heteromonocyclic group containing
 1 to 4 nitrogen atom(s), carbamoyl(C₁-C₄)alkoxy substituted
 15 with saturated 3 to 8-membered heteromonocyclic group
 containing 1 to 4 nitrogen atom(s) having aryl(lower)alkyl,
 phenylcarbamoyl(C₁-C₄)alkoxy having (C₇-C₁₄)alkyl,
 phenylcarbamoyl(C₁-C₄)alkoxy having (C₁-C₄)alkyl,
 phenylcarbamoyl(C₁-C₄)alkoxy having tri-halo(C₁-C₄)alkyl,
 20 phenylcarbamoyl(C₁-C₄)alkoxy having nitro,
 phenylcarbamoyl(C₁-C₄)alkoxy having halogen,
 phenylcarbamoyl(C₁-C₄)alkoxy having di(C₁-C₄)alkylamino-
 (C₁-C₄)alkyl, carbonyl(C₁-C₄)alkoxy substituted with
 saturated 3 to 8-membered heteromonocyclic group containing 1
 25 to 4 nitrogen atom(s) having benzhydryl, (C₁-C₄)alkyl-
 carbamoyl(C₁-C₄)alkoxy having benzhydryl, carbamoyl(C₁-C₄)-
 alkoxy substituted with unsaturated 3 to 8-membered
 heteromonocyclic group containing 1 to 2 sulfur atom(s) and 1
 to 3 nitrogen atom(s) having (C₁-C₄)alkyl substituted with
 30 saturated 3 to 8-membered heteromonocyclic group containing 1
 to 4 nitrogen atom(s) substituted with unsaturated condensed
 heterocyclic group containing 1 to 4 nitrogen atom(s),
 carbonyl(C₁-C₄)alkoxy having unsaturated 3 to 8-membered
 heteromonocyclic group containing 1 to 4 nitrogen atom(s)
 35 substituted with oxo and phenyl, N-(C₁-C₄)alkyl-phenyl-

(C₁-C₄)alkylcarbamoyl(C₁-C₄)alkoxy, N-phenyl(C₁-C₄)alkyl-
 phenyl(C₁-C₄)alkylcarbamoyl(C₁-C₄)alkoxy, N-(C₁-C₄)alkyl-
 (C₁-C₄)alkylcarbamoyl(C₁-C₄)alkoxy having unsaturated 3 to 8-
 5 membered heteromonocyclic group containing 1 to 4 nitrogen
 atom(s), N-(C₁-C₄)alkyl-(C₁-C₄)alkylcarbamoyl(C₁-C₄)alkoxy
 having unsaturated condensed heterocyclic group containing 1
 to 4 nitrogen atom(s), phenylcarbamoyl(C₁-C₄)alkoxy having
 (C₁-C₄)alkyl, phenylcarbamoyl(C₁-C₄)alkoxy having
 (C₁-C₄)alkoxy, phenylcarbamoyl(C₁-C₄)alkoxy having halogen,
 10 carbamoyl(C₁-C₄)alkoxy having unsaturated 3 to 8-membered
 heteromonocyclic group containing 1 to 2 sulfur atom(s) and 1
 to 3 nitrogen atom(s), carbonyl(C₁-C₄)alkoxy substituted with
 unsaturated condensed heterocyclic group containing 1 to 4
 nitrogen atom(s),
 15 phenylcarbamoyl(C₁-C₄)alkoxy having di-halo(C₁-C₄)alkoxy,
 phenylcarbamoyl(C₁-C₄)alkoxy having di(C₁-C₄)alkylamino,
 N-phenyl-phenyl(C₁-C₄)alkylcarbamoyl(C₁-C₄)alkoxy,
 N-(C₁-C₄)alkyl-cyclo(C₄-C₆)alkylcarbamoyl(C₁-C₄)alkoxy,
 and the most preferred one may be carbamoylmethoxy,
 20 ethylcarbamoylmethoxy, sec-butylcarbamoylmethoxy,
 n-butylcarbamoylmethoxy, hexylcarbamoylmethoxy,
 cyclohexylcarbamoylmethoxy, phenylcarbamoylmethoxy,
 tetrazolylcarbamoylmethoxy, indolylcarbamoylmethoxy,
 N-methyl-methylcarbamoylmethoxy,
 25 N-methyl-butylcarbamoylmethoxy, benzylcarbamoylmethoxy,
 methoxyethylcarbamoylmethoxy,
 methylthioethylcarbamoylmethoxy,
 dimethylaminoethylcarbamoylmethoxy,
 piperidinocarbonylmethoxy, guanidinocarbonylmethoxy,
 30 hydroxybutylcarbamoylmethoxy, 1-methylbutylcarbamoylmethoxy,
 2-hydroxypropylcarbamoylmethoxy,
 ethoxypropylcarbamoylmethoxy, hydroxypropylcarbamoylmethoxy,
 diethylaminopropylcarbamoylmethoxy,
 tri-fluoroethylcarbamoylmethoxy,
 35 1-ethoxycarbonyl-ethoxycarbonylmethylcarbamoylmethoxy,

morpholinoethylcarbamoylmethoxy,
 pyridylethylcarbamoylmethoxy, pyridylmethylcarbamoylmethoxy,
 imidazolylpropylcarbamoylmethoxy,
 phenylbutylcarbamoylmethoxy, carbamoylmethoxy having
 5 piperidyl substituted with benzyl,
 octylphenylcarbamoylmethoxy, butylphenylcarbamoylmethoxy,
 tri-fluoromethylphenylcarbamoylmethoxy,
 nitrophenylcarbamoylmethoxy, fluorophenylcarbamoylmethoxy,
 dimethylaminomethylbenzylcarbamoylmethoxy,
 10 piperadinocarbonylmethoxy substituted with benzhydryl,
 benzhydrylmethylcarbamoylmethoxy, carbamoylmethoxy having
 thiazolyl having piperidinomethyl substituted with indolyl,
 piperidinocarbonylmethoxy having piridazinyl substituted with
 phenyl and oxo, N-methyl-benzylcarbamoylmethoxy,
 15 N-benzyl-benzylcarbamoylmethoxy,
 N-methoxy-indolyethylcarbamoylmethoxy,
 N-methyl-pyridylethylcarbamoylmethoxy, tolylcarbamoylmethoxy,
 methoxyphenylcarbamoylmethoxy, chlorophenylcarbamoylmethoxy,
 thiazolylcarbamoylmethoxy, dihydroindolylcarbonylmethoxy,
 20 di-fluoromethoxyphenylcarbamoylmethoxy,
 N-ethyl-propylcarbamoylmethoxy,
 N-methyl-methylpropylcarbamoylmethoxy,
 N-ethyl-butylcarbamoylmethoxy, N-methyl-
 hexylcarbamoylmethoxy, fluorophenylcarbamoylmethoxy,
 25 dimethylaminophenylcarbamoylmethoxy,
 N-phenyl-benzylcarbamoylmethoxy,
 N-methyl-phenethylcarbamoylmethoxy and
 N-methyl-cyclohexylcarbamoylmethoxy.

30 Suitable example of "lower alkyl" in R^5 can be referred
 to aforementioned "lower alkyl", in which the preferred one
 may be (C_1-C_4) alkyl, and the most preferred one may be
 methyl.

35 Suitable example of "aryl" in R^5 can be referred to

aforementioned "aryl", in which the preferred one may be phenyl.

Suitable example of "lower alkylene" moiety in the term of "lower alkylene which may have one or more suitable substituent(s)" may include straight or branched one such as methylene, ethylene, trimethylene, 1-methylethylene, tetramethylene, pentamethylene, hexamethylene, methylmethylene, 1-ethylethylene, 2-ethylpropylene, and the like, in which the preferred one may be (C₁-C₄)alkylene, and the most preferred one may be methylene, ethylene and 1-methylethylene.

Suitable example of "suitable substituent(s)" moiety in the term of "lower alkylene which may have one or more suitable substituent(s)" may include aryl, in which the preferred one may be phenyl.

Suitable example of "lower alkenylene" may include straight or branched one having 2 to 6 carbon atom(s) such as vinylene, propenylene, butenylene, 1 or 2 or 3-pentenylene, 1 or 2 or 3-hexenylene, methylvinylene, ethylvinylene, 1 or 2 or 3-methylpropenylene, 1 or 2 or 3-ethylpropenylene, 1 or 2 or 3 or 4-methyl-1 or 2-butenylene, and the like, in which the preferred one may be (C₂-C₄)alkenylene, and the most preferred one may be vinylene.

The processes for preparing the object compound [I] are explained in detail in the following.

Process 1

The object compound [I] or a salt thereof can be prepared by reacting a compound [II] with a compound [III] or a salt thereof.

Suitable salt of the compound [III] may be the same as those exemplified for the compound [I].

The reaction is preferably carried out in the presence

of a base such as an alkali metal carbonate [e.g. sodium carbonate, potassium carbonate, etc.], an alkaline earth metal carbonate [e.g. magnesium carbonate, calcium carbonate, etc.], an alkali metal bicarbonate [e.g. sodium bicarbonate, potassium bicarbonate, etc.], tri(lower)alkylamine [e.g. trimethylamine, triethylamine, etc.], picoline or the like.

The reaction is usually carried out in a conventional solvent, such as an alcohol [e.g. methanol, ethanol, propanol, isopropanol, etc.], diethyl ether, tetrahydrofuran, dioxane, or any other organic solvent which does not adversely influence the reaction.

The reaction temperature is not critical, and the reaction can be carried out under cooling to heating.

Process 2

The object compound [Ib] or a salt thereof can be prepared by subjecting a compound [Ia] or a salt thereof to elimination reaction of the amino protective group.

Suitable salts of the compounds [Ia] may be the same as those exemplified for the compound [I].

This reaction is carried out in accordance with a conventional method such as hydrolysis, reduction or the like.

The hydrolysis is preferably carried out in the presence of a base or an acid including Lewis acid.

Suitable base may include an inorganic base and an organic base such as an alkali metal [e.g. sodium, potassium, etc.], an alkaline earth metal [e.g. magnesium, calcium, etc.], the hydroxide or carbonate or bicarbonate thereof, hydrazine, trialkylamine [e.g. trimethylamine, triethylamine, etc.], picoline, 1,5-diazabicyclo[4.3.0]non-5-ene, 1,4-diazabicyclo[2.2.2]octane, 1,8-diazabicyclo[5.4.0]undec-7-ene, or the like.

Suitable acid may include an organic acid [e.g. formic acid, acetic acid, propionic acid, trichloroacetic acid,

trifluoroacetic acid, etc.], an inorganic acid [e.g. hydrochloric acid, hydrobromic acid, sulfuric acid, hydrogen chloride, hydrogen bromide, hydrogen fluoride, etc.] and an acid addition salt compound [e.g. pyridine hydrochloride, etc.].

The elimination using trihaloacetic acid [e.g. trichloroacetic acid, trifluoroacetic acid, etc.] or the like is preferably carried out in the presence of cation trapping agents [e.g. anisole, phenol, etc.].

The reaction is usually carried out in a solvent such as water, an alcohol [e.g. methanol, ethanol, etc.], methylene chloride, chloroform, tetrachloromethane, tetrahydrofuran, a mixture thereof or any other solvent which does not adversely influence the reaction. A liquid base or acid can be also used as the solvent. The reaction temperature is not critical and the reaction is usually carried out under cooling to heating.

The reduction method applicable for the elimination reaction may include chemical reduction and catalytic reduction.

Suitable reducing agents to be used in chemical reduction are a combination of metal [e.g. tin, zinc, iron, etc.] or metallic compound [e.g. chromium chloride, chromium acetate, etc.] and an organic or inorganic acid [e.g. formic acid, acetic acid, propionic acid, trifluoroacetic acid, p-toluenesulfonic acid, hydrochloric acid, hydrobromic acid, etc.].

Suitable catalysts to be used in catalytic reduction are conventional ones such as platinum catalysts [e.g. platinum plate, spongy platinum, platinum black, colloidal platinum, platinum oxide, platinum wire, etc.], palladium catalysts [e.g. spongy palladium, palladium black, palladium oxide, palladium on carbon, colloidal palladium, palladium on barium sulfate, palladium on barium carbonate, etc.], nickel catalysts [e.g. reduced nickel, nickel oxide, Raney nickel,

etc.], cobalt catalysts [e.g. reduced cobalt, Raney cobalt, etc.], iron catalysts [e.g. reduced iron, Raney iron, etc.], copper catalysts [e.g. reduced copper, Raney copper, Ullman copper, etc.] and the like.

5 In case that the amino protective group is benzyl, the reduction is preferably carried out in the presence of a combination of palladium catalysts [e.g. palladium black, palladium on carbon, etc.] and formic acid or its salt [e.g. ammonium formate, etc.].


10 The reduction is usually carried out in a conventional solvent which does not adversely influence the reaction such as water, an alcohol [e.g. methanol, ethanol, propanol, etc.], chlorobenzene, N,N-dimethylformamide, or a mixture thereof. Additionally, in case that the above-mentioned
15 acids to be used in chemical reduction are in liquid, they can also be used as a solvent. Further, a suitable solvent to be used in catalytic reduction may be the above-mentioned solvent, and other conventional solvent such as diethyl ether, dioxane, tetrahydrofuran, etc. or a mixture thereof.

20 The reaction temperature of this reduction is not critical and the reaction is usually carried out under cooling to heating.

25 The object compound [I] or a salt thereof possesses gut selective sympathomimetic, anti-ulcerous, anti-pancreatitis, lipolytic and anti-pollakiuria activities, and are useful for the treatment and/or prevention of gastrointestinal disorders caused by smooth muscle contractions in human beings or animals, and more particularly to methods for the treatment
30 and/or prevention of spasm or hyperanakisia in case of irritable bowel syndrome, gastritis, gastric ulcer, duodenal ulcer, enteritis, cholecystopathy, cholangitis, urinary calculus and the like; for the treatment and/or prevention of ulcer such as gastric ulcer, duodenal ulcer, peptic ulcer,
35 ulcer causes by non steroidal anti-inflammatory drugs, or the

like; for the treatment and/or prevention of dysuria such as pollakiuria, urinary incontinence or the like in case of nervous pollakiuria, neurogenic bladder dysfunction, nocturia, unstable bladder, cystospasm, chronic cystitis, chronic prostatitis, overflow incontinence, passive incontinence, reflux incontinence, urge incontinence, urinary incontinence, or the like; and for the treatment and/or prevention of pancreatitis, obesity, diabetes, glycosuria, hyperlipidemia, hypertension, atherosclerosis, glaucoma, melancholia, depression, and the like.

The following compound to be used as the β_3 adrenergic receptor agonist which is shown by formulas (IV), (V), (VI), (VII) and (VIII) are also useful for the therapeutic treatment of dysuria, and the like.



or a salt thereof, wherein


A is a bond, $-(CH_2)_n-$ or $CH(B)-$, where n is an integer of 1 to 3 and

B is $-CN$, $-CON(R^9)R^{9'}$ or $-CO_2R^7$;

R^1 is lower alkyl, aryl or arylalkyl;

R^2 is hydrogen, hydroxy, alkoxy, $-CH_2OH$, cyano, $-C(O)OR^7$, $-CO_2H$, $-CONH_2$, tetrazole, $-CH_2NH_2$ or halogen;

R^3 is hydrogen, alkyl, heterocycle or



5

R⁴ is hydrogen, alkyl or B;

R⁵, R^{5'}, R⁸, R^{8'} and R^{8''} are independently hydrogen,

alkoxy, lower alkyl, halogen, -OH, -CN,

10

-(CH₂)_nNR⁶COR⁷, -CON(R⁶)R^{6'}, -CON(R⁶)OR^{6'}, -CO₂R⁶,

-SR⁷, -SOR⁷, -SO₂R⁷, -N(R⁶)SO₂R¹, -N(R⁶)R^{6'},

-NR⁶COR⁷, -OCH₂CON(R⁶)R^{6'}, -OCH₂CO₂R⁷ or aryl; or

R⁵ and R^{5'} or R⁸ and R^{8'} may together with the carbon atoms to which they are attached form an aryl or heterocycle;

15

R⁶ and R^{6'} are independently hydrogen or lower alkyl; and

R⁷ is lower alkyl;

R⁹ and R^{9'} are independently hydrogen, lower alkyl,

20

alkyl, cycloalkyl, arylalkyl, aryl, heteroaryl; or

R⁹ and R^{9'} may together with the nitrogen atom to which they are attached form a heterocycle;

with the proviso that when


A is a bond or -(CH₂)_n and

25

R³ is hydrogen or unsubstituted alkyl,

then R⁴ is B or substituted alkyl:

30



35 or a salt thereof, wherein

n is 0 to 5;

m is 0 or 1;

r is 0 to 3;

A is pyridinyl;

5 R^1 is (1) hydroxy, (2) oxo, (3) halogen, (4) cyano,
 (5) NR^8R^8 , (6) SR^8 , (7) trifluoromethyl,
 (8) C_1-C_{10} alkyl, (9) OR^8 , (10) SO_2R^9 , (11) $OCOR^9$,
 (12) NR^8COR^9 , (13) COR^9 , (14) $NR^8SO_2R^9$,
 (15) $NR^8CO_2R^8$, or (16) C_1-C_{10} alkyl substituted by
 10 hydroxy, halogen, cyano, NR^8R^8 , SR^8 ,
 trifluoromethyl, OR^8 , C_3-C_8 cycloalkyl, phenyl,
 NR^8COR^9 , COR^9 , SO_2R^9 , $OCOR^9$, $NR^8SO_2R^9$ or $NR^8CO_2R^8$;

R^2 and R^3 are independently (1) hydrogen, (2) C_1-C_{10}
 alkyl or (3) C_1-C_{10} alkyl with 1 to 4 substituents
 15 selected from hydroxy, C_1-C_{10} alkoxy, and halogen;

X is (1) $-CH_2-$, (2) $-CH_2-CH_2-$, (3) $-CH=CH-$ or
 (4) $-CH_2O-$;

R^4 and R^5 are independently (1) hydrogen,
 (2) C_1-C_{10} alkyl, (3) halogen, (4) NHR^8 , (5) OR^8 ,
 20 (6) SO_2R^9 or (7) $NHSO_2R^9$;

R^6 is (1) hydrogen or (2) C_1-C_{10} alkyl;

R^7 is $Z-(R^{1a})_n$;

R^{1a} is (1) R^1 , (2) C_3-C_8 cycloalkyl, (3) phenyl
 optionally substituted with up to 4 groups
 25 independently selected from R^8 , NR^8R^8 , OR^8 , SR^8 and
 halogen, or (4) 5 or 6-membered heterocycle with
 from 1 to 4 heteroatoms selected from oxygen,
 sulfur and nitrogen, optionally substituted with up
 to four groups independently selected from oxo, R^8 ,
 30 NR^8R^8 , OR^8 , SR^8 , and halogen;

Z is (1) phenyl, (2) naphthyl, (3) a 5 or 6-membered
 heterocyclic ring with from 1 to 4 heteroatoms
 selected from oxygen, sulfur and nitrogen, (4) a
 benzene ring fused to a C_3-C_8 cycloalkyl ring,
 35 (5) a benzene ring fused to a 5 or 6-membered

heterocyclic ring with from 1 to 4 heteroatoms selected from oxygen, sulfur and nitrogen,


(6) a 5 or 6-membered heterocyclic ring with from 1 to 4 heteroatoms selected from oxygen, sulfur and nitrogen fused to a 5 or 6-membered heterocyclic ring with from 1 to 4 heteroatoms selected from oxygen, sulfur and nitrogen, or

(7) a 5 or 6-membered heterocyclic ring with from 1 to 4 heteroatoms selected from oxygen, sulfur and nitrogen fused to a C₃-C₈ cycloalkyl ring;

R⁸ is (1) hydrogen, (2) C₁-C₁₀ alkyl, (3) C₃-C₈ cycloalkyl, (4) Z optionally having 1 to 4 substituents selected from halogen, nitro, oxo, NR¹⁰R¹⁰, C₁-C₁₀ alkyl, C₁-C₁₀ alkoxy, C₁-C₁₀ alkylthio, and C₁-C₁₀ alkyl having 1 to 4 substituents selected from hydroxy, halogen, CO₂H, CO₂-C₁-C₁₀ alkyl, SO₂-C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl, C₁-C₁₀ alkoxy, and Z optionally substituted by from 1 to 3 of halogen, C₁-C₁₀ alkyl or C₁-C₁₀ alkoxy, or (5) C₁-C₁₀ alkyl having 1 to 4 substituents selected from hydroxy, halogen, CO₂H, CO₂-C₁-C₁₀ alkyl, SO₂-C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl, C₁-C₁₀ alkoxy, C₁-C₁₀ alkyl, and Z optionally substituted by from 1 to 4 of halogen, C₁-C₁₀ alkyl or C₁-C₁₀ alkoxy;

R⁹ is (1) R⁸ or (2) NR⁸R⁸; and


R¹⁰ is (1) C₁-C₁₀ alkyl, or (2) two R¹⁰ groups together with the N to which they are attached formed a 5 or 6-membered ring optionally substituted with C₁-C₁₀ alkyl:



or a salt thereof, wherein

X is hydrogen, halogen, trifluoromethyl or lower alkyl,
and

R is hydrogen; lower alkyl which may have a suitable
substituent selected from the group consisting of
cyclo(C₃-C₇)alkyl, hydroxy, lower alkoxy, carboxy
and lower alkoxy carbonyl; cyclo(C₃-C₇)alkyl or
lower alkanoyl:



or a salt, ester or amide thereof, wherein

R¹ is a hydrogen, fluorine, chlorine or bromine atom or
a hydroxyl, hydroxymethyl, methyl, methoxyl, amino,
formamido, acetamido, methylsulphonylamido, nitro,
benzyloxy, methylsulphonylmethyl, ureido,
trifluoromethyl or p-methoxybenzylamino group;

R² is a hydrogen, fluorine, chlorine or bromine atom or
a hydroxyl group;

R³ is a hydrogen, chlorine or bromine atom or a hydroxyl
group,

R⁴ is a hydrogen atom or a methyl group;


R⁵ is a hydrogen atom or a methyl group;

R^6 is a hydrogen, fluorine or chlorine atom or a methyl, methoxyl or hydroxy group;

X is an oxygen atom or a bond;

Y is an alkylene group of up to 6 carbon atoms or a bond; and

Z is an alkylene, alkenylene or alkynylene group of up to 10 carbon atoms: or



(VIII)

or a salt thereof, wherein

R is hydrogen or methyl,

R^1 is hydrogen, halogen, hydroxy, benzyloxy, amino or hydroxymethyl,

R^2 is hydrogen, hydroxymethyl, $-NHR^3$, $-SO_2NR^4R^{4'}$ or nitro,

R^3 is hydrogen, methyl, $-SO_2R^5$, formyl or $-CONHR^{6'}$,

R^4 and $R^{4'}$ are independently hydrogen, lower alkyl or benzyl,

R^5 is lower alkyl, benzyl or $-NR^4R^{4'}$,

R^6 is hydrogen or lower alkyl,

$R^{6'}$ is hydrogen or lower alkyl, and,

X is N, O, S or methylene;

when X is N, O or S,

then R^9 is hydrogen,

either R^7 or R^8 is hydrogen, and

the other is hydrogen, amino, acetylamino or hydroxy;

when X is methylene,

then both R^7 and R^8 is hydrogen, and

R^9 is hydrogen, amino, acetylamino or hydroxy.

The compounds (IV), (V), (VI), (VII) and (VIII) identified above or a salt thereof to be used in the present invention are the known ones and disclosed in EP-A2-659737, U.S. Patent 5,561,142, EP-A1-211721, EP-A1-023385, and WO97/25311, respectively.

In the formula (IV), the terms "alk" or "alkyl" refers to both straight and branched chain groups having 1 to 12 carbon atoms, preferably 1 to 8 carbons. It is understood, therefore, that the terms "alk" and "alkyl" denote both unsubstituted groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl, 1-methylbutyl, 2,2-dimethylbutyl, 2-methylpentyl, n-hexyl and the like as well as substituted groups. The term "substituted alkyl" specifically denotes an alkyl group as defined above having one or more of the following substituents: halo (especially to form trihaloalkyl, particularly trichloromethyl or trifluoromethyl); aryl; cycloalkyl; hydroxy; amino; thiol; or Y, where Y is -CN, alkoxy, $-\text{CON}(\text{R}^6)\text{R}^{6'}$, $-\text{CO}_2\text{R}^6$ or $-\text{N}(\text{R}^6)\text{SO}_2\text{R}^1$.

The term "lower alkyl" as employed herein includes such alkyl groups as described above containing 1 to 6 carbon atoms.

The term "alkoxy" refers to any of the above alkyl groups linked to an oxygen atom.

The term "lower alkoxy" refers to any of the above lower alkyl groups linked to an oxygen atom.

The term "aryl" refers to monocyclic or bicyclic aromatic groups containing from 6 to 10 carbons in the ring portion, such as phenyl, naphthyl, or such groups optionally substituted with one or more substituents selected from hydrogen, alkoxy, lower alkyl, halogen, -OH, -CN, $-(\text{CH}_2)_n\text{NR}^6\text{COR}^7$, $-\text{CON}(\text{R}^6)\text{R}^{6'}$, $-\text{CON}(\text{R}^6)\text{OR}^{6'}$, $-\text{CO}_2\text{R}^6$, $-\text{SOR}^7$,

$-\text{SO}_2\text{R}^7$, $-\text{N}(\text{R}^6)\text{SO}_2\text{R}^1$, $-\text{N}(\text{R}^6)\text{R}^{6'}$, $-\text{NR}^6\text{COR}^7$, $-\text{OCH}_2\text{CON}(\text{R}^6)\text{R}^{6'}$, $-\text{OCH}_2\text{CO}_2\text{R}^7$ or aryl. Phenyl and substituted phenyl are preferred.

The term "halogen" or "halo" refers to chlorine, bromine, fluorine or iodine.

The term "heterocycle" refers to fully saturated or unsaturated rings of 5 or 6 atoms containing one or two oxygen and/or sulfur atoms and/or one to four nitrogen atoms provided that the total number of heteroatoms in the ring is four or less. Preferred monocyclic heterocycle groups include 2- and 3-thienyl, 2- and 3-furyl, 2-, 3- and 4-pyridyl and imidazolyl. The term heterocycle also includes bicyclic rings wherein the five- or six-membered ring containing oxygen and/or sulfur and/or nitrogen atoms as defined above is fused to a benzene ring and the bicyclic ring is attached by way of an available carbon atom. Preferred bicyclic heterocycle groups include 4-, 5-, 6- or 7-indolyl, 4-, 5-, 6- or 7-isoindolyl, 5-, 6-, 7- or 8-quinolinyl, 5-, 6-, 7- or 8-isoquinolinyl, 4-, 5-, 6- or 7-benzothiazolyl, 4-, 5-, 6- or 7-benzoxazolyl, 4-, 5-, 6- or 7-benzimidazolyl, 4-, 5-, 6- or 7-benzoxadiazolyl, 4-, 5-, 6- or 7-benzofuranzanyl, 4-, 5-, 6- or 7-benzodioxolyl and 4-, 5-, 6- or 7-benzofuran. The term "heterocycle" also includes such monocyclic and bicyclic rings wherein an available carbon atom is substituted with one or more substituents selected from nitro, keto, azo, thiazo, hydrogen, alkoxy, lower alkyl, halogen, $-\text{OH}$, $-\text{CN}$, $-(\text{CH}_2)_n\text{NR}^6\text{COR}^7$, $-\text{CON}(\text{R}^6)\text{R}^{6'}$, $-\text{CON}(\text{R}^6)\text{OR}^{6'}$, $-\text{CO}_2\text{R}^6$, $-\text{SR}^7$, $-\text{SOR}^7$, $-\text{SO}_2\text{R}^7$, $-\text{N}(\text{R}^6)\text{SO}_2\text{R}^1$, $-\text{N}(\text{R}^6)\text{R}^{6'}$, $-\text{NR}^6\text{COR}^7$, $-\text{OCH}_2\text{CON}(\text{R}^6)\text{R}^{6'}$, $-\text{OCH}_2\text{CO}_2\text{R}^7$ or aryl.

In the formula (V), the alkyl groups specified above are intended to include those alkyl groups of the designated length in either a straight or branched configuration. Exemplary of such alkyl groups are methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, tertiary butyl, pentyl,

isopentyl, hexyl, isohexyl, and the like.

The alkoxy groups specified above are intended to include those alkoxy groups of the designated length in either a straight or branched configuration. Exemplary of
 5 such alkoxy groups are methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, tertiary butoxy, pentoxy, isopentoxy, hexoxy, isohexoxy and the like.

The term "halogen" is intended to include the halogen atoms fluorine, chlorine, bromine and iodine.

10 Examples of 5 and 6-membered heterocycles and fused heterocycles of A, Z and R^{1a} include pyridyl, quinolinyl, pyrimidinyl, pyrrolyl, thienyl, imidazolyl, thiazolyl, benzimidazolyl, thiadiazolyl, benzothiadiazolyl, indolyl, indolinyl, benzodioxolyl, benzodioxanyl, benzothiophenyl,
 15 benzofuranyl, benzoxazinyl, benzisoxazolyl, benzothiazolyl, tetrahydronaphthyl, dihydrobenzofuranyl, tetrahydroquinolinyl, furopyridine and thienopyridine.

The preferred values of A and Z are phenyl, naphthyl, benzene ring fused to a 5 or 6-membered heterocyclic ring
 20 with from 1 to 4 heteroatoms selected from oxygen, sulfur and nitrogen, or heterocycles with from 1 to 4 heteroatoms independently selected from one of oxygen or sulfur, and/or 1 to 4 nitrogen atoms.

The more preferred values of A are phenyl, pyridyl, quinolinyl, pyrimidinyl, pyrrolyl, thienyl, imidazolyl, and
 25 thiazolyl.

The more preferred values of Z are phenyl, naphthyl, quinolinyl, thienyl, benzimidazolyl, thiadiazolyl, benzothiadiazolyl, indolyl, indolinyl, benzodioxolyl,
 30 benzodioxanyl, benzothiophenyl, benzofuranyl, benzoxazinyl, benzisoxazolyl, benzothiazolyl, tetrahydronaphthyl, dihydrobenzofuranyl, triazolyl, tetrazolyl, oxadiazolyl, imidazolyl, oxazolyl, thiazolyl, imidazolidinyl, pyrazolyl, isoxazolyl, pyridyl, pyrimidyl, pyrazolyl,
 35 tetrahydrobenzothiazolyl and tetrahydroquinolinyl. When Z is

attached to $-\text{NSO}_2(\text{CH}_2)_r-$, it is preferably phenyl, naphthyl or a benzene ring fused to a 5 or 6-membered heterocyclic ring with from 1 to 4 heteroatoms selected from oxygen, sulfur and nitrogen. When Z is part of the definition of R^8 , it is preferably phenyl, a 5 or 6-membered heterocyclic ring with from 1 to 4 heteroatoms selected from oxygen, sulfur and nitrogen, a benzene ring fused to a 5 or 6-membered heterocyclic ring with from 1 to 4 heteroatoms selected from oxygen, sulfur and nitrogen, or a 5 or 6-membered heterocyclic ring with from 1 to 4 heteroatoms selected from oxygen, sulfur and nitrogen fused to a C_3 - C_8 cycloalkyl ring.

The preferred heterocycles of R^{1a} are thienyl, thiadiazolyl, thiazolyl, tetrazolyl, oxadiazolyl, imidazolyl, oxazolyl, thiazolyl, imidazolidinyl, isoxazolyl, pyridyl, pyrimidyl, and pyrazolyl.

Certain of the above defined terms may occur more than once in the above formula and upon such occurrence each term shall be defined independently of the other; thus for example. NR^8R^8 may represent NH_2 , NHCH_3 , $\text{N}(\text{CH}_3)\text{CH}_2\text{CH}_3$, and the like.

In the formula (VI), the term "lower alkyl" may include straight or branched one having 1 to 6 carbon atom(s), such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, 3-pentyl, isopentyl, tert-pentyl, neopentyl, hexyl, isohexyl, and the like.

The term "halogen" may include fluorine, chlorine, bromine, iodine, and the like.

The term "lower alkoxy" may include methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, t-butoxy, pentyloxy, t-pentyloxy, hexyloxy, and the like.

The term "lower alkoxy carbonyl" may include methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, t-butoxycarbonyl, t-pentyloxycarbonyl, heptyloxycarbonyl, and the like.

The term "lower alkanoyl" may include acetyl, propanoyl, butanoyl, 2-methylpropanoyl, pentanoyl, 2,2-dimethylpropanoyl, hexanoyl, and the like.

5 The term "cyclo(C₃-C₇)alkyl" may include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and the like.

10 In the formula (VII), the term "alkylene" may include methylene, ethylene, propylene, isopropylene, butylene, isobutylene, sec-butylene, tert-butylene, pentylene, tert-pentylene, neo-pentylene, hexylene, iso-hexylene, heptylene, octylene, nonylene, decylene, and the like.

15 The term "alkenylene" may include vinylene, propenylene, butenylene, 1 or 2 or 3-pentenylene, 1 or 2 or 3-hexenylene, methylvinylene, ethylvinylene, heptenylene, octenylene, nonenylene, decenylene, and the like.

20 The term "alkynylene" may include ethynylene, propynylene, butynylene, 1 or 2 or 3-pentynylene, 1 or 2 or 3-hexynylene, heptynylene, octynylene, nonynylene, decynylene, and the like.

25 In the formula (VIII), the term "lower alkyl" may include straight or branched one having 1 to 6 carbon atom(s), such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, 3-pentyl, isopentyl, tert-pentyl, neopentyl, hexyl, isohexyl, and the like.

30 In order to show the usefulness of the compounds (IV) to (VIII) for the prophylactic and therapeutic treatment of above-mentioned diseases in a human being or an animal, the pharmacological test data of the representative compounds thereof are shown in the following.

Test 1

35 Effect on the increase in intravesical pressure induced

by carbachol in anesthetized dog

Test Compound

(1) N-[5-[2-[1-(3,4-Dimethoxyphenyl)-2-phenylethylamino]-
1(R)-hydroxyethyl]-2-hydroxyphenyl]methanesulfonamide
(This compound was obtained according to a similar
manner to that of Example 1 in EP-A2-659737.)

(2) N-[4-[2-[2(R)-Hydroxy-2-(pyridin-3-yl)ethylamino]ethyl]-
phenyl]-4-[4-(3-cyclopentylpropyl)-5-tetrazolon-1-yl]-
benzenesulfonamide
(This compound was obtained according to a similar
manner to that of Example 70 in U.S. Patent 5,561,142.)

(3) N-[5-[2-[2-(9H-Carbazol-2-yloxy)ethylamino]-1-
hydroxyethyl]-2-hydroxyphenyl]methanesulfonamide
hydrochloride
(This compound was obtained according to a similar
manner to that of Example 2 in WO97/25311.)

Test Method

Female Beagle dogs weighing 8.0-15.0 kg were fasted for
24 hours and maintained under halothane anesthesia. A 12F
Foley catheter was lubricated with water soluble jelly,
inserted into the urethral orifice and advanced approximately
10 cm until the balloon tip was placed well inside the
bladder. The balloon was then inflated with 5 ml of room air
and catheter slowly withdrawn just past the first resistance
that is felt at the bladder neck. Urine was completely
drained out through the catheter, and 30 ml of biological
saline was infused. The catheter was connected to pressure
transducer, and intravesical pressure was continuously
recorded. The test compound was injected intravenously at 5
minutes before the administration of carbachol (1.8 µg/kg).

Test Results

| Treatment | Increase in intravesical pressure (mmHg) |
|-----------------------------------|--|
| Control | 5.0±0.9 |
| Test Compound (1) (0.01 mg/kg) | 1.8±0.3** |
| **P<0.01 vs Control (ANOVA) (N=3) | |

| Treatment | Increase in intravesical pressure (mmHg) |
|-----------------------------------|--|
| Control | 5.0±1.2 |
| Test Compound (2) (0.01 mg/kg) | 2.3±1.5* |
| *P<0.05 vs Control (ANOVA) (N=4) | |

| Treatment | Increase in intravesical pressure (mmHg) |
|-----------------------------------|--|
| Control | 5.0±0.5 |
| Test Compound (3) (0.01 mg/kg) | 1.7±0.6** |
| **P<0.01 vs Control (ANOVA) (N=3) | |

The test compounds (1), (2) and (3) inhibited the increase in intravesical pressure induced by carbachol.

Test 2

Effect on cystometrogram in anesthetized rat

Test Compound

(4) 2(S)-[(7-Ethoxycarbonylmethoxy-1,2,3,4-tetrahydro-naphthalen-2-yl)amino]-1(R)-(3-chlorophenyl)ethanol hydrochloride

(This compound was obtained according to a similar

manner to that of Example 11 in EP-A1-211721.)

Test Method

Male S.D. rats, aged 7-9 weeks, were anesthetized with urethane (1.2 g/kg, sc) and supine on a board. After a midline incision in the abdomen, the bladder was exposed. Polyethylene catheters, which were used to measure the intravesical pressure and to infuse saline, were inserted into the bladder through a small incision at the apex of the bladder dome. The bladder catheters were connected to an infusion pump and pressure transducer respectively. Saline was infused at a rate of 5.0 ml/hr for 1 hour. Cystometrography was performed at 15 minutes intervals for 1 hour. The bladder capacity (volume) was calculated from the time required to fill the bladder. The test compound was administered intravenously via femoral vein.

Test Result

| Time (minutes) | Vehicle | compound (4) (0.01 mg/kg) |
|------------------|------------------------------------|---------------------------|
| | (relative % of before value) (N=5) | |
| Before (Control) | 100 | 100 |
| 0-15 | 128 | 152 |
| 15-30 | 111 | 146 |
| 30-45 | 114 | 143 |
| 45-60 | 104 | 151 |

The test compound (4) increased in bladder capacity at a dose of 0.01 mg/kg.

Test 3

Effect on rhythmic contraction of bladder in anesthetized rat

Test Compound

(5) Methyl 4-[2(R)-[2-(3-chlorophenyl)-2(R)-hydroxyethyl-amino]propyl]phenoxyacetate

(This compound was obtained according to a similar manner to that of Example 6 in EP-A1-023385.)

Test Method

Male S.D. rats, aged 7-8 weeks, were anesthetized with urethane (1.2 g/kg, ip) and supine on a board. After a midline incision in the abdomen, the bladder was exposed. Balloon catheter, which was used to measure the intravesical pressure, was inserted into the bladder through a small incision at the apex of the bladder dome. The balloon was then inflated with distilled water and the balloon port of the catheter was connected to pressure transducer. Rhythmic contraction of bladder induced by increase intravesical pressure to 10 mmHg. The test compound was administered intravenously via femoral vein.

Test Result

| | | compound (5) (0.1 mg/kg) | |
|--|------------------|------------------------------|-------|
| | | (relative % of before value) | (N=3) |
| Time (minutes) | | | |
| 25 | Before (Control) | 100 | |
| | 30 | 66.3** | |
| **P<0.01 vs Before value (paired t-test) | | | |

The test compound (5) decreased in contractile force of bladder at a dose of 0.1 mg/kg.

Suitable salt of the compounds in the present invention is a pharmaceutically acceptable and conventional non-toxic salt, and may include a salt with a base or an acid addition salt such as a salt with an inorganic base, for example, an

alkali metal salt (e.g., sodium salt, potassium salt, etc.),
an alkaline earth metal salt (e.g., calcium salt, magnesium
salt, etc.), an ammonium salt;

a salt with an organic base, for example, an organic amine
5 salt (e.g., triethylamine salt, pyridine salt, picoline salt,
ethanolamine salt, triethanolamine salt, dicyclohexylamine
salt, N,N'-dibenzylethylenediamine salt, etc.);

an inorganic acid addition salt (e.g., hydrochloride,
hydrobromide, sulfate, phosphate, etc.);

10 an organic carboxylic sulfonic acid addition salt (e.g.,
formate, acetate, trifluoroacetate, maleate, tartrate,
fumarate, methanesulfonate, benzenesulfonate,
toluenesulfonate, etc.);

a salt with a basic or acidic amino acid (e.g., arginine,
15 aspartic acid, glutamic acid, etc.).

The compounds in the present invention can be isolated
and purified by a conventional method such as pulverization,
recrystallization, column-chromatography, high-performance
20 liquid chromatography (HPLC), reprecipitation, desalting
resin column chromatography, or the like.

The compound in the present invention may be obtained as
its hydrate, and its hydrate is included within the scope of
present invention.

25 It is to be noted that each of the compounds in the
present invention may include one or more stereoisomer such
as optical isomer(s) and geometrical isomer(s) due to
asymmetric carbon atom(s) and double bond(s) and all such
isomers and the mixture thereof are included within the scope
30 of the present invention.

The compounds in the present invention or a salt thereof
include solvated compound [e.g., enclosure compound (e.g.,
hydrate, etc.)].

35 The compounds in the present invention or a salt thereof
include both its crystal form and non-crystal form.

It should be understood that the compounds in the present invention may include the prodrug form.

This application is based on applications No. PP2826/98 and PP5058/98 filed in Australia, the content of which is
5 incorporated hereinto by reference.

The compounds in the present invention may be obtained by methods exemplified by EP-A2-659737, U.S. Patent 5,561,142, EP-A1-211721, EP-A1-023385, or WO97/25311.

10 The patent, patent applications and publications cited herein are incorporated by reference.

The pharmaceutical composition of the present invention can be used in the form of a pharmaceutical preparation, for
15 example, in solid, semisolid or liquid form, which contains a compound in the present application or a pharmaceutically acceptable salt thereof, as an active ingredient in admixture with an organic or inorganic carrier or excipient suitable for rectal, pulmonary (nasal or buccal inhalation), nasal,
20 ocular, external (topical), oral or parenteral (including subcutaneous, intravenous and intramuscular) administrations or insufflation.

The active ingredient may be compounded, for example,
25 with the usual non-toxic, pharmaceutically acceptable carriers for tablets, pellets, troches, capsules, suppositories, creams, ointments, aerosols, powders for insufflation, solutions, emulsions, suspensions, and any other form suitable for use. And, if necessary, in addition,
30 auxiliary, stabilizing, thickening and coloring agents and perfumes may be used.

The compound or a salt thereof in the present invention is included in the pharmaceutical composition in an amount
35 sufficient to produce the desired effect upon the process or

condition of the diseases.

The pharmaceutical composition of the present invention can be manufactured by the conventional method in this field of the art. If necessary, the technique generally used in this field of the art for improving the bioavailability of a drug can be applied to the pharmaceutical composition of the present invention.

For applying the composition to a human being or an animal, it is preferable to apply it by intravenous (including i.v. infusion), intramuscular, pulmonary, or oral administration, or insufflation.

While the dosage of therapeutically effective amount of the compound in the present invention varies from and also depends upon the age and condition of each individual patient to be treated, in the case of intravenous administration, a daily dose of 0.01-100 mg of the compound in the present invention per kg weight of a human being or an animal, in the case of intramuscular administration, a daily dose of 0.01-100 mg of the compound in the present invention per kg weight of a human being or an animal, in case of oral administration, a daily dose of 0.01-200 mg of the compound in the present invention per kg weight of a human being or an animal is generally given for the prophylactic and/or therapeutic treatment of above-mentioned diseases in a human being or an animal.

The following Preparations and Examples are given for the purpose of illustrating this invention.

Preparation 1

Under nitrogen, a solution of N-benzyl-(3-ethoxycarbonylmethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (2.0 g) in tetrahydrofuran (20 ml) was added
 5 di-tert-butyl dicarbonate (1.4 g) at 5°C, and the solution was stirred at room temperature for 2.5 hours. The resulting solution was evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to
 10 afford [8-(N-benzyl-tert-butoxycarbonylamino)-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid ethyl ester (2.58 g).

NMR (CDCl₃, δ) : 1.29 (3H, t, J=7.1Hz), 1.08-1.63 (11H, m), 1.87-2.18 (3H, m), 2.50-2.70 (3H, m), 3.17-3.35 (1H, m), 4.27 (2H, q, J=7.1Hz), 4.37-4.70 (2H, m),
 15 4.55 (2H, s), 6.60 (1H, dd, J=2.5 and 8.1Hz), 6.93 (1H, d, J=8.3Hz), 7.19-7.83 (4H, m)

Preparation 2

To a solution of [8-(N-benzyl-tert-butoxycarbonylamino)-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid ethyl ester (2.5 g) in ethanol (25 ml) was added 1N sodium hydroxide aqueous solution at 5°C, and the solution was stirred at room temperature for 1 hour. The resulting solution was evaporated in vacuo to afford sodium [8-(N-benzyl-tert-butoxycarbonylamino)-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetate (2.35 g).
 25

NMR (CDCl₃, δ) : 1.04-1.60 (10H, m), 1.75-2.11 (3H, m), 2.41-2.67 (4H, m), 3.15-3.55 (1H, m), 4.05 (2H, s), 4.27-4.65 (2H, m), 6.52 (2H, d, J=7.9Hz), 6.88 (1H, d, J=8.2Hz), 7.12-7.42 (5H, m)
 30

Preparation 3

Under nitrogen, to a mixture of sodium [8-(N-benzyl-tert-butoxycarbonylamino)-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetate (0.50 g) in dichloromethane
 35

(5 ml) were added 4N hydrogenchloride in ethyl acetate (0.28 ml), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.24 g), aniline (112 μ l) and a catalytic amount of 4-dimethylaminopyridine at 5°C, and the mixture was stirred at room temperature for 3 hours. The resulting mixture was poured into 1N hydrogen chloride aqueous solution and extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and brine, dried over magnesium sulfate, and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford N-benzyl-N-tert-butoxycarbonyl-(3-N-phenylcarbamoylmethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (0.47 g).

NMR (CDCl_3 , δ) : 1.13-1.75 (11H, m), 1.90-2.25 (3H, m), 2.50-2.80 (3H, m), 3.24-3.38 (1H, m), 4.27-4.78 (2H, m), 4.53 (2H, s), 6.68 (1H, dd, $J=2.5$ and 8.1Hz), 6.98 (1H, d, $J=8.3\text{Hz}$), 7.12-7.43 (9H, m), 7.58-7.65 (2H, m)

The following compound was obtained according to a similar manner to that of Preparation 3.

Preparation 4

N-Benzyl-N-tert-butoxycarbonyl-(3-N-butylcarbamoylmethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine

NMR (CDCl_3 , δ) : 0.93 (3H, t, $J=7.2\text{Hz}$), 1.10-2.20 (17H, m), 2.50-2.75 (3H, m), 3.15-3.45 (4H, m), 4.28-4.77 (2H, m), 4.40 (2H, s), 6.60 (2H, dd, $J=2.4$ and 8.1Hz), 6.95 (1H, d, $J=8.3\text{Hz}$), 7.25-7.50 (5H, m)

Preparation 5

Under nitrogen, 4N hydrogenchloride in ethyl acetate (4 ml) was added to N-benzyl-N-tert-butoxycarbonyl-(3-N-phenylcarbamoylmethoxy-6,7,8,9-tetrahydro-5H-

benzocyclohepten-6-yl)amine (0.45 g) at room temperature, and the solution was stirred at the same temperature for 3.5 hours. The resulting solution was evaporated in vacuo. The residue was dissolved in a mixture of saturated aqueous sodium bicarbonate solution and ethyl acetate. After separation, the organic layer was dried over magnesium sulfate and evaporated in vacuo to afford 2-(8-benzylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy)-N-phenylacetamide (346 mg).

NMR (CDCl_3 , δ) : 1.38-1.62 (1H, m), 1.63-2.14 (4H, m), 2.65-2.82 (3H, m), 2.90-2.98 (2H, m), 3.82 (2H, ABq, $J=3.7$ and 12.7Hz), 4.58 (2H, s), 6.71 (1H, dd, $J=2.7$ and 8.2Hz), 6.82 (1H, d, $J=2.7\text{Hz}$), 7.04 (1H, d, $J=8.2\text{Hz}$), 7.13-7.42 (8H, m), 7.56-7.65 (2H, m)

MASS (m/z) : 401 ($M+H$)⁺

The following compound was obtained according to a similar manner to that of Preparation 5.

Preparation 6

2-(8-Benzylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy)-N-butylacetamide

NMR (CDCl_3 , δ) : 0.92 (3H, t, $J=7.2\text{Hz}$), 1.25-1.61 (4H, m), 1.73-2.30 (4H, m), 2.63-3.05 (5H, m), 3.34 (2H, q, $J=6.9\text{Hz}$), 3.85 (2H, ABq, $J=13.1$ and 17.0Hz), 4.45 (2H, s), 6.64 (1H, dd, $J=2.7$ and 8.2Hz), 7.01 (1H, d, $J=8.2\text{Hz}$), 7.20-7.45 (5H, m)

MASS (m/z) : 381 ($M+H$)⁺

Preparation 7

A solution of N-benzyl-(6,7,8,9-tetrahydro-2-hydroxy-5H-benzocyclohepten-6-yl)amine (194 mg) and di-tert-butyl dicarbonate (174 mg) in tetrahydrofuran (2 ml) was stirred at room temperature for 20 hours and partitioned between ethyl acetate and an aqueous solution of sodium bicarbonate. The

organic layer was separated, washed with brine, dried over sodium sulfate, and evaporated in vacuo. The residue was chromatographed (toluene - ethyl acetate) over silica gel (5.8 g) to afford N-benzyl-N-(6,7,8,9-tetrahydro-2-hydroxy-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine as a pale yellow solid (196 mg).

mp : 150-156°C (dec.)

IR (KBr) : 3348 (br), 1666, 1246 cm^{-1}

NMR (DMSO- d_6 , δ) : 1.2-1.6 (10H, m), 1.96 (3H, m), 2.49-2.7 (3H, m), 3.10-3.85 (2H, m), 4.30-4.7 (2H, m), 6.46-6.54 (2H, m), 6.77 (1H, m), 7.15-7.36 (5H, m)

(+) API-ES MASS (m/z) : 390 (M^+ +Na)

Preparation 8

A solution of N-benzyl-N-(6,7,8,9-tetrahydro-2-hydroxy-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine (166 mg) and N-chlorosuccinimide (66.4 mg) in 1,4-dioxane (2 ml) was refluxed for 28 hours, cooled to room temperature, and partitioned between ethyl acetate and brine. The organic layer was separated, washed with brine, dried over sodium sulfate, and evaporated in vacuo. The residue was chromatographed (toluene -ethyl acetate) over silica gel (3.6 g) to afford N-benzyl-N-(3-chloro-6,7,8,9-tetrahydro-2-hydroxy-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine (139 mg) as a colorless gum.

IR (KBr) : 3411 (br), 3263 (br), 1655, 1252 cm^{-1}

Preparation 9

A mixture of N-benzyl-N-(3-chloro-6,7,8,9-tetrahydro-2-hydroxy-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine (135 mg) and 4N hydrogen chloride in ethyl acetate (3 ml) was stirred under ice-cooling for 2 hours and at room temperature for 5 hours and evaporated in vacuo. The residue was partitioned between ethyl acetate and an aqueous solution of

sodium bicarbonate. The organic layer was separated, washed with brine, dried over sodium sulfate, and evaporated in vacuo to afford N-benzyl-(3-chloro-6,7,8,9-tetrahydro-2-hydroxy-5H-benzocyclohepten-6-yl)amine (100 mg) as a yellow gum.

IR (KBr) : 3419 (br), 1238 cm^{-1}

NMR (CDCl_3 , δ) : 1.48-1.59 (1H, m), 1.71-2.03 (3H, m), 2.3-3.16 (6H, m), 3.78 (1H, d, $J=13.0\text{Hz}$), 3.87 (1H, d, $J=13.0\text{Hz}$), 6.73 (1H, s), 7.08 (1H, s), 7.15-7.32 (6H, m)

Preparation 10

Under nitrogen, to a suspension of sodium hydride (60% in oil, 47 mg) in N,N-dimethylformamide (3 ml) was dropwise added 8-benzylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ol (300 mg) in N,N-dimethylformamide (2 ml) at 5°C, and the mixture was stirred at the same temperature for 30 minutes. To this one was added n-propyl bromide (0.10 ml), and the mixture was stirred at the same temperature for 3 hours. The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate, and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 50:1 to 30:1) to give N-benzyl-(3-isopropoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (278 mg).

(+) ESI-MASS (m/z) : 310 ($\text{M}+\text{H}$)⁺

The following compounds [Preparation 11 and 12] were obtained according to a similar manner to that of Preparation 10.

Preparation 11

N-Benzyl-(3-hexyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine

(+) APCI-MASS (m/z) : 352 (M+H)⁺

Preparation 12

N-Benzyl-(3-benzyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine

NMR (CDCl₃, δ) : 1.2-1.6 (2H, m), 1.7-2.1 (3H, m), 2.6-2.95 (5H, m), 3.7-3.9 (2H, m), 5.03 (2H, s), 6.70 (1H, dd, J=2.7 and 8.2Hz), 6.83 (1H, d, J=2.6Hz), 6.98 (1H, d, J=8.2Hz), 7.2-7.5 (10H, m)

Preparation 13

Under nitrogen, a solution of (2S)-1-[N-benzyl-(3-benzyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol (20 g), chlorotriethylsilane (7.4 ml), imidazole (3.2 g) and 4-dimethylaminopyridine (0.48 g) in N,N-dimethylformamide (200 ml) was stirred at room temperature for 12 hours. The resulting mixture was poured into water, and extracted with ethyl acetate-hexane (1:1). The organic layer was successively washed with saturated aqueous sodium hydrogencarbonate and brine, dried over anhydrous magnesium sulfate, and evaporated in vacuo to give N-benzyl-N-(3-benzyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-[(2S)-3-phenoxy-2-(triethylsilyloxy)-propyl]amine (25 g).

(+) ESI-MASS (m/z) : 622 (M+H)⁺

Preparation 14

A mixture of N-benzyl-N-(3-benzyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]amine (25 g) and 10% palladium on activated carbon (50% wet, 2.5 g) in methanol (250 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 5 hours. After filtration, the filtrate was evaporated in vacuo to give 8-[(2S)-3-phenoxy-2-(triethylsilyloxy)propylamino]-6,7,8,9-tetrahydro-5H-

benzocyclohepten-2-ol (17 g).

(+) APCI-MASS (m/z) : 442 (M+H)⁺

Preparation 15

Under nitrogen, a mixture of 8-[(2S)-3-phenoxy-2-(triethylsilyloxy)propylamino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ol (17 g) and di-tert-butyl dicarbonate (9.5 g) in tetrahydrofuran (180 ml) was stirred at room temperature for 18 hours. After evaporation, the residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 20:1 to 10:1) to give N-(3-hydroxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-butoxycarbonylamine (17 g).

(+) APCI-MASS (m/z) : 442 (M-Boc+2H)⁺

Preparation 16

Under nitrogen, to a suspension of sodium hydride (60% in oil, 12 mg) in N,N-dimethylformamide (2 ml) was dropwise added N-(3-hydroxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-butoxycarbonylamine (150 mg) in N,N-dimethylformamide (1 ml) at 5°C, and the mixture was stirred at the same temperature for 30 minutes. To this one was added ethyl iodide (27 μ l), and the mixture was stirred at room temperature for 2 hours. The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate, and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 20:1) to give N-(3-ethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-butoxycarbonylamine (130 mg).

NMR (CDCl₃, δ) : 0.55-0.75 (6H, m), 0.9-1.0 (9H, m),
1.2-1.55 (4H, m), 1.48 (9H, m), 1.9-2.2 (3H, m),
2.55-2.9 (3H, m), 3.1-3.65 (4H, m), 3.85-4.1 (4H,

m), 4.2-4.4 (1H, m), 6.55-6.7 (2H, m), 6.85-7.0
(4H, m), 7.2-7.3 (2H, m)
(+) ESI-MASS (m/z) : 592 (M+Na)⁺

5 The following compound was obtained according to a
similar manner to that of Preparation 16.

Preparation 17

10 N-(3-Allyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-
yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-
butoxycarbonylamine

15 NMR (DMSO-d₆, δ) : 0.55-0.75 (6H, m), 0.8-1.1 (9H, m),
1.2-1.6 (1H, m), 1.48 (9H, m), 1.9-2.2 (3H, m),
2.6-2.9 (2H, m), 3.15-3.7 (4H, m), 3.8-4.05 (2H,
m), 4.25-4.5 (3H, m), 5.25-5.45 (2H, m), 5.95-6.15
(1H, m), 6.6-6.75 (2H, m), 6.8-7.0 (4H, m), 7.25-
7.3 (2H, m)

Preparation 18

20 Under nitrogen, to dimethylsulfoxide (3 ml) was added
potassium hydroxide (31 mg) at room temperature, and the
suspension was stirred at the same temperature for 1 hour.
To this one were added N-(3-hydroxy-6,7,8,9-tetrahydro-5H-
benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-
25 (triethylsilyloxy)propyl]-tert-butoxycarbonylamine (150 mg)
and isopropyl bromide (78 μl), and the mixture was stirred at
room temperature for 3 hours. The resulting mixture was
poured into aqueous 10% sodium hydrogencarbonate, and
extracted with ethyl acetate. The organic layer was washed
30 with brine, dried over anhydrous magnesium sulfate, and
evaporated in vacuo. The residue was purified by column
chromatography on silica gel (hexane:ethyl acetate = 20:1) to
give N-(3-isopropoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-
6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-
35 butoxycarbonylamine (96 mg).

(+) APCI-MASS (m/z) : 326 (M-Boc+2H)⁺

The following compounds [Preparations 19 to 21] were obtained according to a similar manner to that of Preparation 18.

Preparation 19

N-(3-Benzoyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-butoxycarbonylamine

(+) APCI-MASS (m/z) : 532 (M-Boc+2H)⁺

Preparation 20

N-Benzyl-N-[3-(3-cyanopyridin-2-yloxy)-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]-tert-butoxycarbonylamine

(+) APCI-MASS (m/z) : 370 (M-Boc+2H)⁺

Preparation 21

N-Benzyl-N-(2-chloro-3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine

(+) APCI-MASS (m/z) : 316, 318 (M-Boc+2H)⁺

Preparation 22

Under nitrogen, to a solution of N-(3-hydroxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-butoxycarbonylamine (150 mg) in tetrahydrofuran (5 ml) was added triphenylphosphine (290 mg), cyclopentanol (60 μ l) and diethyl azodicarboxylate (174 μ l) at 5°C, and the mixture was stirred at the same temperature for 5.5 hours. The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate, and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 10:1) to

give N-(3-cyclopentyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-butoxycarbonylamine (108 mg).

(+) APCI-MASS (m/z) : 510 (M-Boc+2H)⁺

5

The following compound was obtained according to a similar manner to that of Preparation 22.

Preparation 23

10 N-(3-Phenethyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-butoxycarbonylamine
(+) APCI-MASS (m/z) : 546 (M-Boc+2H)⁺

15 Preparation 24

Under nitrogen, to a suspension of potassium carbonate (76 mg) in a mixture of acetonitrile (5 ml) and N,N-dimethylformamide (2 ml) was added N-(3-hydroxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-butoxycarbonylamine (150 mg)
20 at room temperature. After being stirred at the same temperature for 30 minutes, to this one was an excessive amount of bromofluoromethane, and the mixture was stirred for 12 hours. The resulting solution was poured into saturated
25 aqueous sodium hydrogencarbonate, and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate =10:1) to give N-(3-fluoromethoxy-
30 6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-butoxycarbonylamine (130 mg).

(+) APCI-MASS (m/z) : 474 (M-Boc+2H)⁺

35 Preparation 25

Under nitrogen, to a solution of N-(3-hydroxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-butoxycarbonylamine (150 mg) in dichloromethane (5 ml) were added pyridine (64 ml),
 5 4-nitrophenyl chloroformate (116 mg) and a catalytic amount of 4-dimethylaminopyridine at 5°C. After being stirred at the same temperature for 2 hours, to this one was added about 6.8M ammonia in ethanol (0.5 ml), and the mixture was stirred at room temperature for 12 hours. The resulting mixture was
 10 pured into water, and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 3:1) to give N-[(2S)-3-phenoxy-2-(triethyl-
 15 silyloxy)propyl]-N-[3-carbamoyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]-tert-butoxycarbonylamine (129 mg).

(+) APCI-MASS (m/z) : 485 (M-Boc+2H)⁺

Preparation 26

20 Under nitrogen, a mixture of 8-benzylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ol (7.7 g) and di-tert-butyl dicarbonate (6.9 g) in tetrahydrofuran (80 ml) was stirred at room temperature for 12 hours. After evaporation, the residue was purified by column chromatography on silica
 25 gel (hexane:ethyl acetate = 7:1) to give N-benzyl-N-(3-hydroxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine (10 g).

NMR (CDCl₃, δ) : 1.1-1.7 (10H, m), 1.85-2.1 (3H, m),
 2.45-2.7 (3H, m), 3.15-3.85 (2H, m), 4.3-4.7 (2H,
 30 m), 5.05-5.25 (1H, m), 6.3-6.6 (2H, m), 6.86 (2H,
 d, J=8.1Hz), 7.15-7.4 (6H, m)

Preparation 27

To N-benzyl-N-[3-(3-cyanopyridin-2-yloxy)-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]-tert-butoxycarbonylamine
 35

(480 mg) was added 4N hydrogen chloride in ethyl acetate (5 ml) at room temperature, and the mixture was stirred at the same temperature for 1 hour. After evaporation in vacuo, the residue was dissolved into a mixture of saturated aqueous sodium hydrogencarbonate and ethyl acetate, followed by being made basic with saturated aqueous sodium hydrogencarbonate. After separation, the organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo to give 2-(8-benzylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yl)oxy-3-cyanopyridine (359 mg).

(+) APCI-MASS (m/z) : 370 (M+H)⁺

The following compounds [Preparations 28 to 30] were obtained according to a similar manner to that of Preparation 27.

Preparation 28

N-Benzyl-[3-(2-methoxyethoxy)-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]amine

(+) APCI-MASS (m/z) : 326 (M+H)⁺

Preparation 29

N-Benzyl-(3-phenyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine

(+) APCI-MASS (m/z) : 328 (M+H)⁺

Preparation 30

N-Benzyl-(2-chloro-3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine

(+) APCI-MASS (m/z) : 316, 318 (M+H)⁺

Preparation 31

Under nitrogen, to dimethylsulfoxide (5 ml) was added potassium hydroxide (34 mg) at room temperature, and the suspension was stirred at the same temperature for 40

minutes. To this one were added N-benzyl-N-(3-hydroxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine (200 mg) and 2-chloroethyl methyl ether (55 μ l), and the mixture was stirred at room temperature for 4 hours. The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 10:1 to 5:1) to give N-benzyl-N-[3-(2-methoxyethoxy)-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]-tert-butoxycarbonylamine (213 mg).

(+) APCI-MASS (m/z) : 326 (M-Boc+2H)⁺

Preparation 32

A solution of 2-(8-benzylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy)acetic acid ethyl ester (320 mg) and 28% ammonium hydroxide (1 ml) in methanol (3 ml) was stirred at room temperature for 12 hours. The mixture was evaporated in vacuo to give 2-(8-benzylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy)acetamide (270 mg).

(+) APCI-MASS (m/z) : 325 (M+H)⁺

Preparation 33

Under nitrogen, to a suspension of sodium hydride (60% in oil, 300 mg) in N,N-dimethylformamide (20 ml) was added N-benzyl-N-(3-hydroxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine (2.5 g) in N,N-dimethylformamide (25 ml) at 5°C. After being stirred at room temperature for 30 minutes, to this one were added ethyl bromoacetate (0.75 ml) and tetra-n-butylammonium bromide (1.1 g) at 5°C, and the mixture was stirred at room temperature for 32 hours. The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous

magnesium sulfate, and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 5:1) to give 2-[8-(N-benzyl-N-tert-butoxycarbonylamino)-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid ethyl ester (1.6 g).

NMR (CDCl₃, δ) : 1.1-1.7 (10H, m), 1.30 (3H, t, J=7.1Hz), 1.85-2.2 (3H, m), 2.5-2.7 (3H, m), 3.2-3.4 (1H, m), 4.27 (2H, q, J=7.2Hz), 4.3-4.7 (2H, m), 4.54 (2H, s), 6.4-6.65 (2H, m), 6.93 (1H, d, J=8.3Hz), 7.2-7.4 (5H, m)

(+) APCI-MASS (m/z) : 354 (M-Boc+2H)⁺

Preparation 34

To a solution of 2-[8-(N-benzyl-N-tert-butoxycarbonylamino)-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid ethyl ester (1.6 g) in ethanol (16 ml) was added aqueous 1N sodium hydroxide at 5°C, and the mixture was stirred at room temperature for 1 hour. After evaporation in vacuo, the residue was dissolved into a mixture of aqueous 0.1N hydrochloric acid and ethyl acetate. After separation, the organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was triturated with hexane and dried in vacuo to give 2-[8-(N-benzyl-N-tert-butoxycarbonylamino)-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid (1.3 g).

NMR (DMSO-d₆, δ) : 1.1-1.7 (10H, m), 1.75-2.2 (3H, m), 2.45-2.7 (1H, m), 3.15-3.75 (4H, m), 4.3-4.7 (2H, m), 4.57 (2H, s), 6.4-6.7 (2H, m), 6.96 (1H, d, J=8.2Hz), 7.2-7.5 (5H, m), 12.9 (1H, br s)

(+) APCI-MASS (m/z) : 326 (M-Boc+2H)⁺

Preparation 35

Under nitrogen, a solution of 2-[8-(N-benzyl-N-tert-butoxycarbonylamino)-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid (300 mg) in dichloromethane (3 ml) were

added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (150 mg), dimethylamine hydrochloride (63 mg) and 4-dimethylaminopyridine (130 mg) at 5°C, and the mixture was stirred at room temperature for 12 hours. The resulting mixture was poured into 0.1N hydrochloric acid and extracted with ethyl acetate. The organic layer was successively washed with saturated aqueous sodium hydrogencarbonate and brine, dried over magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 1:1) to give N-benzyl-N-(3-N,N-dimethylcarbamoylmethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine (312 mg).

NMR (CDCl₃, δ) : 1.1-1.65 (10H, m), 1.85-2.15 (3H, m), 2.5-2.7 (4H, m), 2.98 (3H, s), 3.08 (3H, s), 3.2-3.45 (1H, m), 4.5-4.7 (2H, m), 4.60 (2H, s), 6.45-6.7 (2H, s), 6.93 (1H, d, J=8.2Hz), 7.2-7.4 (5H, m)
(+) APCI-MASS (m/z) : 353 (M-Boc+2H)⁺

Preparation 36

To N-benzyl-N-(3-N,N-dimethylcarbamoylmethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine (279 mg) was added 4N hydrogen chloride in ethyl acetate (3 ml), and the mixture was stirred at room temperature for 1.5 hours. After evaporation in vacuo, the residue was dissolved into saturated aqueous sodium hydrogencarbonate and ethyl acetate, followed by separation. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. A mixture of the residue and 10% palladium on activated carbon (50% wet, 100 mg) in methanol (5 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 7 hours. After filtration, the filtrate was evaporated in vacuo to give 2-(8-amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy)-N,N-dimethylacetamide (137 mg).

NMR (CDCl₃, δ) : 1.4-2.1 (4H, m), 2.65-3.15 (5H, m),

2.98 (3H, m), 3.09 (3H, s), 4.65 (2H, s), 6.68 (1H, dd, J=2.7 and 8.1Hz), 6.77 (1H, d, J=2.7Hz), 6.98 (1H, d, J=8.2Hz)

(+) APCI-MASS (m/z) : 263 (M+H)⁺

5

Preparation 37

Under nitrogen, to a solution of 3-nitro-5,7,8,9-tetrahydrobenzocyclohepten-6-one (300 mg) in 1,2-dichloroethane (10 ml) were added benzylamine (0.24 ml), sodium triacetoxyborohydride (460 mg) and acetic acid (0.17 ml) at room temperature, and the mixture was stirred at the same temperature for 5 hours. The resulting mixture was poured into aqueous 1N sodium hydroxide and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 2:1 to 1:5) to give N-benzyl-(3-nitro-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (342 mg).

10
15
20

(+) APCI-MASS (m/z) : 297 (M+H)⁺

The following compound was obtained according to a similar manner to that of Preparation 37.

25

Preparation 38

N-Benzyl-(6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-amine

(+) APCI-MASS (m/z) : 252 (M+H)⁺

30

Preparation 39

Under nitrogen, to the solution of 1-tetralone (16 g) in dichloromethane (80 ml) were added zinc(II) iodide (0.69 g) and trimethylsilyl cyanide (15 g) at room temperature, and the mixture was stirred at the same temperature for 3 hours.

35

The resulting mixture was poured into saturated aqueous

sodium hydrogencarbonate and extracted with ethyl acetate. The organic layer was successively washed with saturated aqueous sodium hydrogencarbonate and brine, dried over anhydrous magnesium sulfate and evaporated in vacuo to give crude 1-(trimethylsilyloxy)-1-cyano-1,2,3,4-tetrahydronaphthalene (27 g), which was successively used in the next reaction.

The following compounds [Preparations 40 and 41] were obtained according to a similar manner to that of Preparation 39.

Preparation 40

7-Methyl-1-trimethylsilyloxy-1-cyano-1,2,3,4-tetrahydronaphthalene

Preparation 41

7-Chloro-1-trimethylsilyloxy-1-cyano-1,2,3,4-tetrahydronaphthalene

Preparation 42

Under nitrogen, to the suspension of lithium aluminum hydride (8.4 g) in tetrahydrofuran (120 ml) was dropwise added 1-trimethylsilyloxy-1-cyano-1,2,3,4-tetrahydronaphthalene (27 g) in tetrahydrofuran (120 ml) at 5°C, and the mixture was stirred at room temperature for 24 hours. To the mixture were added sodium fluoride (37 g) and water (12 ml) at 5°C, and the mixture was vigorously stirred at room temperature for 30 minutes. The precipitate was removed by filtration. The filtrate was evaporated in vacuo to give 1-aminomethyl-1-hydroxy-1,2,3,4-tetrahydronaphthalene (19 g).

(+) APCI-MASS (m/z) : 178 (M+H)⁺

The following compounds [Preparations 43 and 44] were obtained according to a similar manner to that of Preparation

42.

Preparation 43

1-Aminomethyl-7-chloro-1-hydroxy-1,2,3,4-
5 tetrahydronaphthalene

NMR (CDCl₃, δ) : 1.8-2.2 (4H, m), 2.6-2.95 (4H, m),
7.00 (1H, d, J=8.2Hz), 7.13 (1H, dd, J=2.3 and
8.2Hz), 7.52 (1H, d, J=2.2Hz)

10 Preparation 44

1-Aminomethyl-7-methyl-1-hydroxy-1,2,3,4-
tetrahydronaphthalene

NMR (CDCl₃, δ) : 1.5-2.2 (7H, m), 2.31 (3H, s), 2.6-2.9
15 (4H, m), 6.9-7.05 (2H, m), 7.32 (1H, s)

15 Preparation 45

To a solution of 1-aminomethyl-1-hydroxy-1,2,3,4-
tetrahydronaphthalene (19 g) in 10% acetic acid in water (380
ml) was dropwise added sodium nitrite (11 g) in water (56 ml)
20 at 5°C, and the mixture was stirred at the same temperature
for 2 hours. The resulting mixture was diluted with ethyl
acetate and separated. The organic layer was successively
washed with water, saturated aqueous sodium hydrogencarbonate
and brine, dried over anhydrous magnesium sulfate and
25 evaporated in vacuo. The residue was purified by column
chromatography on silica gel (hexane-ethyl acetate) to give
5,7,8,9-tetrahydrobenzocyclohepten-6-one (9.3 g).

(+) APCI-MASS (m/z) : 161 (M+H)⁺

30 The following compounds [Preparations 46 and 47] were
obtained according to a similar manner to that of Preparation
45.

Preparation 46

35 3-Chloro-5,7,8,9-tetrahydrobenzocyclohepten-6-one

NMR (CDCl_3 , δ) : 1.9-2.1 (2H, m), 2.57 (2H, t, $J=7.0\text{Hz}$), 2.9-3.0 (2H, m), 3.68 (2H, s), 7.09 (1H, d, $J=7.8\text{Hz}$), 7.16 (1H, s), 7.18 (1H, d, $J=7.8\text{Hz}$)

5 Preparation 47

3-Methyl-5,7,8,9-tetrahydrobenzocyclohepten-6-one

NMR (CDCl_3 , δ) : 1.9-2.05 (2H, m), 2.30 (3H, s), 2.57 (2H, t, $J=6.8\text{Hz}$), 2.85-2.95 (2H, m), 3.69 (2H, s), 6.95-7.1 (3H, m)

10 Preparation 48

A stirred solution of (2S)-1-phenoxy-2,3-epoxypropane (1.0 g) and concentrated ammonium hydroxide (10 ml) in ethanol (10 ml) was sealed up at room temperature for 12 hours. The mixture was evaporated in vacuo and dried to give (2S)-1-amino-3-phenoxy-2-propanol (1.1 g).

(+) APCI-MASS (m/z) : 168 (M+H)⁺

20 Preparation 49

To a mixture of N-benzyl-N-(3-hydroxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine (1.01 g), 2,6-lutidine (0.38 ml) and N,N-dimethylaminopyridine (0.067 g) in dichloromethane (20 ml) was added dropwise trifluoromethanesulfonic anhydride (0.51 ml) at -30°C. The mixture was warmed to room temperature over 3 hours and then washed with water, dried over magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel eluting with a mixture of hexane and ethyl acetate (20:1) to afford N-benzyl-N-tert-butoxycarbonyl-(3-trifluoromethylsulfonyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (0.98 g).

NMR ($\text{DMSO}-d_6$, δ) : 1.14-1.50 (11H, m), 1.80-2.15 (3H, m), 2.60-2.80 (3H, m), 3.30-3.60 (1H, m), 4.25-4.62 (2H, m), 6.98-7.50 (8H, m)

(+) ESI-MASS (m/z) : 522 (M+Na)⁺

Preparation 50

To a suspension of N-benzyl-N-tert-butoxycarbonyl-(3-trifluoromethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (500 mg) in toluene (6 ml) and water (2 ml) were added phenylboronic acid (122 mg), sodium carbonate (210 mg) and tetrakis(triphenylphosphine)palladium(0) (35 mg), and the mixture was stirred at 80°C for 3.5 hours. The resulting mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 30:1 to 20:1) to give N-benzyl-N-(3-phenyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine (494 mg).

(+) ESI-MASS (m/z) : 450 (M+Na)⁺

Preparation 51

Under nitrogen, a suspension of N-benzyl-N-(3-hydroxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine (300 mg) and N-chlorosuccinamide (110 mg) in 1,4-dioxane (5 ml) was stirred at 100°C for 9 hours. The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (toluene:ethyl acetate = 50:1 to 10:1) to give N-benzyl-N-(2-chloro-3-hydroxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine (152 mg).

(+) APCI-MASS (m/z) : 302, 304 (M-Boc+2H)⁺

Preparation 52

Under nitrogen, to a solution of (2R)-1-tosyloxy-2,3-epoxypropane (3.0 g) in tetrahydrofuran (30 ml) were added N,N-diisopropylethylamine (2.5 ml) and thiophenol (1.3 ml) at

5°C, and the mixture was stirred at room temperature for 12 hours. The resulting mixture was poured into water and extracted with ethyl acetate. The organic layer was successively washed with saturated aqueous sodium

hydrogencarbonate and brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 5:1 to 3:1) to give (2S)-3-phenylthio-1-tosyloxy-2-propanol (3.9 g).

NMR (CDCl₃, δ) : 2.44 (3H, m), 2.75-3.25 (3H, m), 3.85-4.3 (3H, m), 7.15-7.4 (7H, m), 7.7-7.8 (2H, m)

Preparation 53

Under nitrogen, to a solution of (2S)-3-phenylthio-1-tosyloxy-2-propanol (3.9 g) in ethanol (40 ml) was added 20% sodium methoxide in ethanol (4.7 ml) at 5°C, and the mixture was stirred at the same temperature for 30 minutes. After being filtrated off to remove precipitates, the filtrate was concentrated in vacuo. The residue was dissolved into a mixture of aqueous 0.1N sodium hydroxide and diethyl ether. After separation, the organic layer was successively washed with water and brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 20:1 to 10:1) to give (2S)-1-phenylthio-2,3-epoxypropane (1.5 g).

(+) APCI-MASS (m/z) : 167 (M+H)⁺

Preparation 54

Under nitrogen, to a solution of allylphenylamine (3 ml) in dichloromethane (30 ml) were added pyridine (2.1 ml) and benzyl chloroformate (3.5 ml) at 5°C, and the mixture was stirred at the same temperature for 3 hours. The resulting mixture was poured into aqueous 1N hydrogen chloride and extracted with ethyl acetate. The organic layer was successively washed with saturated aqueous sodium

hydrogencarbonate and brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 20:1 to 10:1) to give N-allyl-N-phenylbenzyloxycarbonylamine (5.5 g).

(+) APCI-MASS (m/z) : 268 (M+H)⁺

Preparation 55

Under nitrogen, to a solution of N-allyl-N-phenylbenzyloxycarbonylamine (5.5 g) in dichloromethane (50 ml) was added m-chloroperbenzoic acid (3.9 g) at 5°C, and the mixture was stirred at room temperature for 3 days. The resulting mixture was poured into aqueous sodium hydrogensulfite and extracted with ethyl acetate. The organic layer was successively washed with saturated aqueous sodium hydrogencarbonate and brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 10:1 to 5:1) to give 1-(N-benzyloxycarbonyl-N-phenylamino)-2,3-epoxypropane (2.2 g).

(+) APCI-MASS (m/z) : 284 (M+H)⁺

Preparation 56

To a suspension of but-3-butenylbenzene (3 ml) and sodium hydrogen carbonate (2.5 g) in a mixture of dichloromethane (200 ml) and water (60 ml) was added small portions of m-chloroperbenzoic acid (3.5 g) at room temperature, and the mixture was stirred at the same temperature for 4 hours. After separation, the organic layer was successively washed with saturated aqueous sodium hydrogencarbonate and brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 100:3) to give 2-phenethyloxirane (970 mg).

NMR (CDCl₃, δ) : 1.7-1.9 (2H, m), 2.45-2.5 (1H, m),

2.7-3.0 (4H, m), 7.1-7.4 (5H, m)

Preparation 57

A mixture of N-benzyl-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (1.0 g) and 10% palladium on activated carbon (50% wet, 300 mg) in methanol (10 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 5.5 hours. After filtration, the filtrate was evaporated in vacuo to give 3-methoxy-(6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (720 mg).

NMR (CDCl₃, δ) : 1.4-1.6 (1H, m), 1.7-2.0 (2H, m), 2.1-2.3 (1H, m), 2.7-2.75 (2H, m), 2.85-3.3 (3H, m), 3.76 (3H, s), 6.66 (1H, dd, J=2.7 and 8.2Hz), 6.76 (1H, d, J=2.6Hz), 7.00 (1H, d, J=8.2Hz)

The following compounds [Preparation 58 and 59] were obtained according to a similar manner to that of Preparation 57.

Preparation 58

2-(8-Amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy)acetic acid tert-butyl ester

NMR (CDCl₃, δ) : 1.3-2.1 (4H, m), 1.49 (9H, s), 2.65-3.1 (5H, m), 4.48 (2H, m), 6.61 (1H, dd, J=2.7 and 8.2Hz), 6.73 (1H, d, J=2.7Hz), 6.98 (1H, d, J=8.2Hz)

Preparation 59

2-(8-Amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy)-N-phenylacetamide

NMR (CDCl₃, δ) : 1.45-2.15 (4H, m), 2.7-3.1 (5H, m), 4.58 (2H, m), 6.72 (1H, dd, J=2.7 and 8.2Hz), 6.83 (1H, d, J=2.7Hz), 7.04 (1H, d, J=8.2Hz), 7.1-7.25 (1H, m), 7.3-7.4 (2H, m), 7.55-7.7 (2H, m), 8.30 (1H, br s)

(+) ESI-MASS (m/z) : 311 (M+H)⁺

Preparation 60

To a solution of N-benzyl-N-(3-hydroxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine (500 mg) in ethyl acetate (5 ml) was added 4N hydrogen chloride in ethyl acetate (5 ml) at room temperature, and the mixture was stirred at the same temperature for 1 hour. The mixture was poured into a mixture of saturated aqueous sodium hydrogencarbonate and ethyl acetate, followed by being made basic with saturated aqueous sodium hydrogencarbonate. After separation, the organic layer was dried over anhydrous magnesium sulfate and evaporated in vacuo to give crude 8-benzylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ol.

Under nitrogen, to a suspension of sodium hydride (60% in oil, 60 mg) in N,N-dimethylformamide (3 ml) was added above obtained one in N,N-dimethylformamide (4 ml) at 5°C. After being stirred at room temperature for 30 minutes, to this one were added tert-butyl bromoacetate (0.22 ml) and tetra-n-butylammonium bromide (220 mg) at 5°C, and the mixture was stirred at room temperature for 6 hours. The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (chloroform:methanol = 50:1 to 20:1) to give 2-(8-benzylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy)acetic acid tert-butyl ester (474 mg).

(+) APCI-MASS (m/z) : 382 (M+H)⁺

Preparation 61

A mixture of 6-(4-nitrophenylazo)pyridin-3-ol (300 mg) and 20% palladium hydroxide on carbon (60 mg) in a mixture of acetic acid (30 ml) and methanol (30 ml) was stirred at room

temperature in the presence of hydrogen at an atmospheric pressure for 70 minutes. After filtration, the filtrate was evaporated in vacuo. Under nitrogen, to a mixture of the residue in dichloromethane (10 ml) was added

5 bis(trimethylsilyl)acetamido (6.0 ml) at 5°C. After being stirred at room temperature for 30 minutes, to this one was added benzyl chloroformate (0.54 ml) at 5°C, and the mixture was stirred at the same temperature for 3 hours. The resulting mixture was poured into saturated aqueous sodium
10 hydrogencarbonate and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. To the residue was added chloroform, and insoluble materials were filtered off. After the filtrate was evaporated in vacuo, the residue
15 was purified by column chromatography on silica gel (chloroform:methanol = 50:1 to 20:1), followed by crystallization from toluene-methanol to give 2-benzyloxy-carbonylamino-5-hydroxypyridine (159 mg).

(+) APCI-MASS (m/z) : 245 (M+H)⁺

20 Preparation 62

Under nitrogen, to a suspension of sodium hydride (60% in oil, 189 mg) in N,N-dimethylformamide (20 ml) was dropwise
25 added 2-benzyloxycarbonylamino-5-hydroxypyridine (1.1 g) in N,N-dimethylformamide (12 ml) at 5°C, and the mixture was stirred at room temperature for 1 hour. To this one was added (2S)-(+)-1-tosyloxy-2,3-epoxypropane (1.1 g) at 5°C, and the mixture was stirred at room temperature for 7 hours. The resulting mixture was poured into saturated aqueous
30 sodium hydrogencarbonate and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (chloroform:ethyl acetate = 9:1) to give 2-benzyloxycarbonyl-
35 amino-5-[(2S)-oxiranylmethoxy]pyridine (780 mg).

(+) APCI-MASS (m/z) : 301 (M+H)⁺

Preparation 63

To a mixture of 8-methoxy-1-tetralone (2.66 g) and zinc iodide (0.096 g) in dichloromethane (13 ml) was dropwise added trimethylsilylcyanide (2.65 ml) under a nitrogen atmosphere. The reaction mixture was stirred at room temperature overnight and then poured into a saturated aqueous sodium bicarbonate solution. The product was extracted with ethyl acetate, and the extract was washed with a saturated aqueous sodium bicarbonate solution and brine, dried over magnesium sulfate, and evaporated in vacuo. The residue was purified by column chromatography on silica gel eluting with a mixture of hexane and ethyl acetate (4:1) to afford 1-cyano-8-methoxy-1-trimethylsilyloxy-1,2,3,4-tetrahydronaphthalene (3.64 g).

NMR (DMSO-d₆, δ) : 0.15 (9H, s), 1.70-1.81 (2H, m), 2.09-2.38 (2H, m), 2.70-2.77 (2H, m), 3.84 (3H, s), 6.76 (1H, d, J=8.0Hz), 6.92 (1H, d, J=8.0Hz), 7.29 (1H, d, J=8.0, 8.0Hz)

(+) ESI-MASS (m/z) : 298 (M+Na)⁺

Preparation 64

To a mixture of lithiumaluminumhydride (0.99 g) in tetrahydrofuran (18 ml) was added dropwise a solution of 1-cyano-8-methoxy-1-trimethylsilyloxy-1,2,3,4-tetrahydronaphthalene (3.59 g) in tetrahydrofuran (18 ml) under ice-cooling and a nitrogen atmosphere. After stirring at room temperature for 3 hours, sodium fluoride (1.09 g) and water (1.41 ml) were added to the reaction mixture under ice-cooling. After stirring for additional 30 minutes, insoluble materials were filtered off and washed with 5% ethanol in ethyl acetate. The filtrate was evaporated in vacuo to afford 1-aminomethyl-1-hydroxy-8-methoxy-1,2,3,4-tetrahydronaphthalene (2.60 g), which was used in the next

step without purification.

NMR (DMSO- d_6 , δ) : 1.38-1.72 (3H, m), 2.07-2.14 (1H, m), 2.62-2.91 (6H, m), 3.77 (3H, s), 6.68 (1H, d, $J=7.8\text{Hz}$), 6.80 (1H, d, $J=7.8\text{Hz}$), 7.09 (1H, dd, $J=7.8, 7.8\text{Hz}$)

(+) ESI-MASS (m/z) : 190 (M-OH)⁺

Preparation 65

To a suspension of 1-aminomethyl-1-hydroxy-8-methoxy-1,2,3,4-tetrahydronaphthalene (2.54 g) in 10% acetic acid was added dropwise a solution of sodium nitrite (1.27 g) in water (6.4 ml) under ice-cooling. The mixture was stirred at the same temperature for 2 hours and then partitioned between ethyl acetate and water. The organic layer was separated, washed with an aqueous sodium bicarbonate solution and brine, dried over magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel eluting with a mixture of hexane and ethyl acetate (10:1) to afford 4-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-one (1.08 g).

NMR (DMSO- d_6 , δ) : 1.77-1.91 (2H, m), 2.47 (2H, t, $J=7.0\text{Hz}$), 2.85-2.92 (2H, m), 3.76 (3H, s), 3.76 (2H, s), 6.76 (1H, d, $J=7.5\text{Hz}$), 6.88 (1H, d, $J=7.5\text{Hz}$), 7.16 (1H, dd, $J=7.5$ and 7.5Hz)

(+) ESI-MASS (m/z) : 213 (M+Na)⁺

Preparation 66

A solution of 4-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-one (500 mg) and benzylamine (0.29 ml) in the presence of catalytic amounts of p-toluenesulfonic acid monohydrate in toluene (5 ml) was refluxed for 2 hours to remove water as the toluene azeotrope, and then the mixture was evaporated in vacuo. To the solution of the residue in methanol (5 ml) was added sodium borohydride (99 mg) under nitrogen at 5°C, and the mixture was stirred at room

temperature for 12 hours. After the resulting mixture was evaporated in vacuo, the residue was dissolved into ice-cold water and ethyl acetate. After separation, the organic layer was washed successively with water and brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 5:1 to 1:1) to give N-benzyl-(4-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (660 mg).

NMR (DMSO- d_6 , δ) : 1.17-1.37 (1H, m), 1.53-2.02 (3H, m), 2.25-2.38 (1H, m), 2.50-3.80 (3H, m), 3.26-3.32 (2H, m), 3.67-3.86 (2H, m), 3.77 (3H, s), 6.68 (1H, d, $J=7.5\text{Hz}$), 6.79 (1H, d, $J=7.5\text{Hz}$), 7.01 (1H, dd, $J=7.5$ and 7.5Hz), 7.17-7.30 (5H, m)

(+) ESI-MASS (m/z) : 282 (M+H)⁺

Preparation 67

Into a mixture of N-benzyl-N-(3-trifluoromethyl-sulfonyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine (0.94 g), palladium acetate (0.084 g), 1,3-bis(diphenylphosphino)propane (0.16 g), triethylamine (0.79 ml) in methanol (2.8 ml) and N,N-dimethylformamide (7 ml) was introduced carbon oxide gas at room temperature for 1 hour. Then the mixture was heated at 75°C for 2.5 hours with carbon oxide gas bubbling. After cooling, the reaction mixture was partitioned between water and ethyl acetate. The organic layer was separated, washed with water (twice) and brine, dried over magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel eluting with a mixture of hexane and ethyl acetate (10:1) to afford 8-(N-benzyl-N-tert-butoxycarbonylamino)-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxylic acid methyl ester (0.76 g).

NMR (DMSO- d_6 , δ) : 1.13-1.50 (11H, m), 1.79-2.13 (3H, m), 2.58-2.79 (3H, m), 3.44-3.74 (1H, m), 3.83 (3H,

s), 4.30-4.62 (2H, m), 7.20-7.38 (6H, m), 7.50-7.60 (1H, m), 7.67 (1H, d, J=7.7Hz)
 (+) ESI-MASS (m/z) : 432 (M+Na)⁺

5 Preparation 68

To a solution of 8-(N-benzyl-N-tert-butoxycarbonylamino)-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxylic acid methyl ester (10.07 g) in dichloromethane (100 ml) was added trifluoroacetic acid (9.47 ml), and the mixture was stirred at room temperature overnight. The mixture was evaporated in vacuo and partitioned between ethyl acetate and an aqueous sodium bicarbonate solution. The organic layer was separated, washed with water and brine, dried over magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel eluting with a mixture of chloroform and methanol (20:1) to afford 8-benzylamino-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxylic acid methyl ester (7.78 g).

NMR (DMSO-d₆, δ) : 1.24-1.43 (1H, m), 1.59-2.08 (4H, m), 2.42-2.50 (1H, m), 2.76-3.03 (4H, m), 3.75 (2H, d, J=3.1Hz), 3.83 (3H, s), 7.16-7.36 (6H, m), 7.67 (1H, dd, J=7.7, 1.8Hz), 7.75 (1H, d, J=1.8Hz)
 (+) ESI-MASS (m/z) : 310 (M+H)⁺

25 Preparation 69

To a mixture of 8-[N-tert-butoxycarbonyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxylic acid methyl ester (4.80 g), imidazole (1.68 g) and catalytic amount of N,N-dimethylaminopyridine in N,N-dimethylformamide (100 ml) was added triethylsilyl chloride (3.82 ml) under ice-cooling. The reaction mixture was stirred at room temperature for 6 hours and partitioned between ethyl acetate and aqueous sodium bicarbonate solution. The organic layer was separated, washed with water (twice) and brine, dried over

magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel eluting with a mixture of hexane and ethyl acetate (20:1) to afford 8-[N-tert-butoxycarbonyl-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]amino]-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxylic acid methyl ester (6.07 g).

NMR (DMSO- d_6 , δ) : 0.50-0.67 (6H, m), 0.83-0.95 (9H, m), 1.13-1.37 (1H, m), 1.42 (9H, s), 1.89-2.22 (3H, m), 2.68-2.98 (3H, m), 3.20-3.48 (4H, m), 3.81, 3.83 (total 3H, s), 3.83-3.92 (1H, m), 3.97-4.07 (1H, m), 4.20-4.37 (1H, m), 6.86-6.97 (3H, m), 7.18-7.34 (3H, m), 7.68-7.75 (2H, m)
(+) APCI-MASS (m/z) : 484 (M-Boc+2H)⁺

Preparation 70

To a cold (-78°C) solution of 8-[N-tert-butoxycarbonyl-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]amino]-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxylic acid methyl ester (5.90 g) in tetrahydrofuran (120 ml) was added dropwise a solution of diisobutylaluminum hydride in hexane (0.95M, 53.2 ml). The reaction mixture was warmed to room temperature over 4 hours, and then saturated sodium bicarbonate solution was added dropwise to the mixture. The resulting precipitate was filtered off and washed with ethyl acetate. The filtrate was extracted with ethyl acetate, and the organic layer was washed with water and brine, dried over magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel eluting with a mixture of hexane and ethyl acetate (5:1) to afford N-(3-hydroxymethyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-butoxycarbonylamine (5.20 g).

NMR (DMSO- d_6 , δ) : 0.52-0.68 (6H, m), 0.85-0.97 (9H, m), 1.08-1.38 (1H, m), 1.41 (9H, s), 1.80-2.20 (3H, m), 2.50-2.88 (3H, m), 3.15-3.52 (4H, m), 3.80-3.90

(1H, m), 3.94-4.05 (1H, m), 4.22-4.40 (1H, m), 4.41
(2H, d, J=4.5Hz), 5.06 (1H, t, J=4.5Hz), 6.83-7.03
(6H, m), 7.30 (2H, dd, J=7.4 and 7.4Hz)
(+) APCI-MASS (m/z) : 456 (M-Boc+2H)⁺

5

Preparation 71

To a mixture of N-(3-hydroxymethyl-6,7,8,9-tetrahydro-
5H-benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyl-
oxy)propyl]-tert-butoxycarbonylamine (1.93 g) and
10 triethylamine (0.58 ml) in dichloromethane (40 ml) was added
dropwise methanesulfonyl chloride (0.27 ml) under ice-
cooling. After stirring at room temperature for 6 hours, the
reaction mixture was washed successively with water, sodium
bicarbonate solution, water and brine. The organic layer was
15 dried over magnesium sulfate and evaporated in vacuo to
afford crude N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-N-
(3-methylsulfonyloxymethyl-6,7,8,9-tetrahydro-5H-
benzocyclohepten-6-yl)-tert-butoxycarbonylamine (2.18 g),
which was used without purification.

20

Preparation 72

To a solution of N-tert-butoxycarbonyl-N-[(2S)-3-
phenoxy-2-(triethylsilyloxy)propyl]-(3-N,N-dimethylamino-
methyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine
25 (0.60 g) in tetrahydrofuran (3 ml) was added a solution of
dimethylamine in tetrahydrofuran (2M, 1.89 ml) under ice-
cooling. The mixture was stirred at room temperature
overnight and then partitioned between water and ethyl
acetate. The organic layer was separated, washed with water
and brine, dried over magnesium sulfate and evaporated in
30 vacuo. The residue was purified by column chromatography on
silica gel eluting with a mixture of chloroform and methanol
(50:1 to 20:1) to afford N-(3-N,N-dimethylaminomethyl-
6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-3-
35 phenoxy-2-(triethylsilyloxy)propyl]-tert-

butoxycarbononylamine (0.41 g).

NMR (DMSO- d_6 , δ) : 0.43-0.67 (6H, m), 0.82-0.98 (9H, m), 1.10-1.35 (1H, m), 1.41 (9H, s), 1.81-2.08 (3H, m), 2.11 (6H, s), 2.50-2.90 (3H, m), 3.17-3.58 (5H, m), 3.78-4.22 (4H, m), 6.87-7.04 (6H, m), 7.22-7.34 (2H, m)

(+) APCI-MASS (m/z) : 583 (M+H)⁺

Preparation 73

A mixture of 8-benzylamino-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxylic acid methyl ester (4.21 g) and sodium methoxide (2.21 g) in formamide (42 ml) was heated at 110°C for 2 hours. After cooling, the reaction mixture was poured into water, and the product was extracted with ethyl acetate. The organic layer was washed with water (twice) and brine, dried over magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel eluting with a mixture of chloroform and methanol (30:1 to 10:1) to afford 8-benzylamino-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxamide (2.71 g).

NMR (DMSO- d_6 , δ) : 1.24-1.42 (1H, m), 1.59-2.05 (4H, m), 2.40-2.50 (1H, m), 2.70-3.01 (4H, m), 3.67-3.84 (2H, m), 7.13 (1H, d, J=7.7Hz), 7.18-7.36 (6H, m), 7.59 (1H, dd, J=7.7 and 1.8Hz), 7.70 (1H, d, J=1.8Hz), 7.86 (1H, s)

(+) APCI-MASS (m/z) : 295 (M+H)⁺

Preparation 74

To a solution of (R)-4-(3-toluenesulfonyloxy-2-hydroxy)propyl-1-benzyloxybenzene (110 mg) in acetonitrile (10 ml), a solution of nitronium tetrafluoroborate (536 mg) in acetonitrile (10 ml) was added dropwise with ice-bath cooling. The reaction mixture was stirred for 0.5 hour and worked up in a usual manner. The crude product was purified by a column chromatography (SiO₂ 50 cm³, eluent; 33% ethyl

acetate-hexane) to give (R)-1-(3-toluenesulfonyloxy-2-hydroxy)propyl-3-nitro-4-benzyloxybenzene (259 mg) as an oil.

IR (Film) : 1623, 1533, 1356, 1267, 1174, 1095, 984,
816 cm^{-1}

5 NMR (CDCl_3 , δ) : 2.17 (1H, d, $J=4.4\text{Hz}$), 2.46 (3H, s),
2.7-2.8 (2H, m), 3.9-4.2 (3H, m), 5.21 (2H, s),
7.06 (1H, t, $J=8.7\text{Hz}$), 7.3-7.5 (7H, m), 7.66 (1H,
d, $J=2.1\text{Hz}$), 7.7-7.9 (3H, m)

10 Preparation 75

To a solution of 4-benzyloxybromobenzene (263 mg) in dry tetrahydrofuran (3 ml), butyllithium (1.54M hexane solution, 0.71 ml) was added dropwise at -78°C under nitrogen, and the resulting mixture was stirred for 30 minutes. To the
15 reaction mixture, boranetrifluoride diethyl ether (123 μl), and a solution of (R)-1-tosyloxy-2,3-epoxypropane (202 mg) in dry tetrahydrofuran (1 ml) were added below -50°C . The reaction mixture was worked up in a usual manner and purified by a column chromatography (SiO_2 40 cm^3 , 25% ethyl acetate-
20 hexane) to give (R)-4-(3-toluenesulfonyloxy-2-hydroxy)propyl-1-benzyloxybenzene (144 mg) as an oil.

IR (Film) : 1512, 1360, 1240, 1178, 978, 833, 814 cm^{-1}

25 NMR (CDCl_3 , δ) : 2.04 (1H, d, $J=4.4\text{Hz}$), 2.45 (3H, s),
2.72 (2H, d, $J=5.3\text{Hz}$), 3.9-4.1 (3H, m), 5.04 (2H,
s), 6.89 (2H, d, $J=8.7\text{Hz}$), 7.06 (2H, d, $J=8.7\text{Hz}$),
7.3-7.5 (7H, m), 7.7-7.9 (3H, m)

Preparation 76

A mixture of (R)-1-(3-toluenesulfonyloxy-2-hydroxy)propyl-3-nitro-4-benzyloxybenzene (26 mg), iron
30 powder, ammonium chloride (2.6 mg) and wet ethanol (0.5 ml) was heated at 80°C for 1 hour and 20 minutes, filtered and worked up in the usual manner. The residue was treated with small amounts of potassium carbonate in methanol and worked
35 up in the usual manner to give epoxide. The crude epoxide

was dissolved in dichloromethane, treated with triethylamine followed by methanesulfonyl chloride, worked up in the usual manner and purified by preparative TLC (SiO₂, 50% ethyl acetate-hexane) to give (R)-1-(3-methanesulfonylamino-4-benzyloxy)phenyl-2,3-epoxypropane (10.8 mg) as an oil.

IR (Film) : 2931, 1612, 1508, 1365, 1321, 1270, 1160, 975, 924 cm⁻¹

NMR (CDCl₃, δ) : 2.50 (1H, dd, J=2.6 and 4.9Hz), 2.7-3.0 (3H, m), 2.89 (3H, s), 3.1-3.2 (1H, m), 5.14 (2H, s), 7.02 (1H, d, J=8.4Hz), 7.24 (1H, dd, J=2.2 and 9.6Hz), 7.3-7.6 (6H, m)

Preparation 77

Under nitrogen, to a solution of 4-acetylamino-phenol (569 mg) and sodium hydride (150 mg) in N,N-dimethylformamide (30 ml) was added (2S)-(+)-1-tosyloxy-2,3-epoxypropane (1.0 g) at 0°C, and the mixture was stirred at the same temperature for 0.5 hour. The mixture was allowed to warm to room temperature and stirred for 3 hours at this temperature. The resulting mixture was poured into 10% aqueous ammonium chloride solution and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford (2S)-1-(4-acetylamino)phenoxy-2,3-epoxypropane (285 mg).

(+) APCI-MASS (m/z) : 149 (M+Na)⁺

Preparation 78

Under nitrogen, to a solution of 2,3-dimethylbenzene-1,4-diol (25.0 g) and potassium hydroxide (40.6 g) in dimethylsulfoxide (100 ml) was added iodomethane (17 ml) at room temperature, and the mixture was stirred for 3 hours at this temperature. The resulting mixture was poured into water and extracted with ethyl acetate. The organic layer

was washed with brine, dried over magnesium sulfate and evaporated in vacuo to afford 1,4-dimethoxy-2,3-dimethylbenzene (23.16 g).

(+) APCI-MASS (m/z) : 167 (M+H)⁺

5

Preparation 79

A solution of 1,4-dimethoxy-2,3-dimethylbenzene (23.16 g), benzoyl peroxide (673 mg) and N-bromosuccinimide (52.0 g) in carbontetrachloride (140 ml) was refluxed for 4 hours and
10 filtered. The filtrate was evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford 2,3-dibromomethyl-1,4-dimethoxybenzene (8.17 g).

(+) APCI-MASS (m/z) : 325 (M+H)⁺

15

Preparation 80

To a solution of N-isopropylcyclohexylamine (3.52 ml) in tetrahydrofuran (10 ml) was added 1M n-butyllithium in hexane (13.8 ml). After being stirred for 20 minutes at this temperature, the solution was treated dropwise with tert-
20 butyl acetate (2.96 ml). And after stirred for 20 minutes, a solution of 2,3-dibromomethyl-1,4-dimethoxybenzene (2.73 g) in tetrahydrofuran (10 ml) was added below -68°C. After all had been introduced, the mixture was allowed to warm to -23°C and stirred for 2.5 hours at this temperature. The resulting
25 mixture was added dropwise to 1N hydrochloric acid (10 ml) with stirring under ice-cooling over 30 minutes and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over
30 silica gel to afford 3-[2-(2-tert-butoxycarbonyl-ethyl)-3,6-dimethoxyphenyl]propionic acid tert-butyl ester (2.69 g).

NMR (CDCl₃, δ) : 1.47 (18H, s), 2.29-2.43 (4H, m),
2.88-2.96 (4H, m), 3.75 (6H, s), 6.66 (2H, s)

35

Preparation 81

3-[2-(2-tert-butoxycarbonylethyl)-3,6-dimethoxyphenyl]-propionic acid tert-butyl ester (6.33 g) was added dropwise to a refluxed solution of sodium hydride (578 mg) in dry toluene (60 ml) and dry tert-butyl alcohol (0.5 ml), and stirred for 3 hours. After cooling, to the reaction mixture glacial acetic acid was added dropwise followed by addition of ice water and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. To the residue were added methanol (50 ml) and 6N hydrochloric acid (15 ml), and this mixture was refluxed for 3 hours, poured into ice and extracted with ethyl acetate. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford 1,4-dimethoxy-5,6,8,9-tetrahydrobenzocyclohepten-7-one (2.99 g).

MASS (m/z) : 221 (M+H)⁺

Preparation 82

A solution of 1,4-dimethoxy-5,6,8,9-tetrahydrobenzocyclohepten-7-one (670 mg) and benzylamine (0.5 ml) in toluene (60 ml) in the presence of a catalytic amount of p-toluenesulfonic acid monohydrate was refluxed for 2 hours to remove water as the toluene azeotrope, and then the mixture was evaporated in vacuo. To the residue in methanol (60 ml) was added sodium borohydride (0.5 g) under nitrogen at 5°C, and the mixture was stirred at room temperature for 12 hours. The resulting mixture was poured into ice-cold water and stirred for 30 minutes before adding ethyl acetate and brine. After separation, the organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate-methanol) over silica gel to afford N-benzyl-(1,4-dimethoxy-5,6,8,9-tetrahydrobenzocyclohepten-7-yl)amine (600 mg).

MASS (m/z) : 312 (M+H)⁺

Preparation 83

Under nitrogen, to fuming nitric acid (60 ml) was added 6,7,8,9-tetrahydrobenzocyclohepten-5-one (20 g) on ice-cooling, and the mixture was stirred at the same temperature for 1 hour. The resulting mixture was poured into ice water (200 ml). The precipitates were collected by filtration and dissolved into ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo to afford 3-nitro-6,7,8,9-tetrahydrobenzocyclohepten-5-one (14.87 g).

(+) APCI-MASS (m/z) : 228 (M+Na)⁺

Preparation 84

A mixture of 3-nitro-6,7,8,9-tetrahydrobenzocyclohepten-5-one (14.87 g), iron (14.87 g) and ammonium chloride (2.0 g) in ethanol (200 ml) and water (40 ml) was refluxed for 5 hours, and filtered. The filtrate was evaporated in vacuo to afford 3-amino-6,7,8,9-tetrahydrobenzocyclohepten-5-one (12.86 g).

NMR (CDCl₃, δ) : 1.75-1.90 (4H, m), 2.60-2.90 (6H, m), 6.73-7.07 (3H, m)

Preparation 85

To a solution of 3-amino-6,7,8,9-tetrahydrobenzocyclohepten-5-one (12.53 g) in 10% sulfuric acid (130 ml) was added sodium nitrite (4.9 g) and sodium sulfite (694 mg) under ice-cooling. After being stirred for 20 minutes at this temperature, to the reaction mixture was added toluene (100 ml), stirred at the room temperature for 18 hours and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo to afford 3-hydroxy-6,7,8,9-tetrahydrobenzocyclohepten-5-one (7.08 g).

(+) APCI-MASS (m/z) : 177 (M+H)⁺

Preparation 86

A mixture of 3-hydroxy-6,7,8,9-tetrahydrobenzocyclohepten-5-one (7.02 g), potassium carbonate (8.25 g) and iodomethane (7.4 ml) in N,N-dimethylformamide (70 ml) was stirred at 50-55°C for 7 hours. The resulting mixture was diluted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo to afford 3-methoxy-6,7,8,9-tetrahydrobenzocyclohepten-5-one (5.49 g).

(+) APCI-MASS (m/z) : 191 (M+H)⁺

Preparation 87

A solution of 3-methoxy-6,7,8,9-tetrahydrobenzocyclohepten-5-one (6.0 g) and benzylamine (3.4 ml) in toluene (60 ml) in the presence of a catalytic amount of p-toluenesulfonic acid monohydrate was refluxed for 2 hours to remove water at the toluene azeotrope, and then the mixture was evaporated in vacuo. To the residue in methanol (60 ml) was added sodium borohydride (1.2 g) under nitrogen at 5°C, and the mixture was stirred at room temperature for 12 hours. The resulting mixture was poured into ice-cold water and stirred for 30 minutes. After separation, the organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate-methanol) over silica gel to afford N-benzyl-(1-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (7.27 g).

(+) APCI-MASS (m/z) : 282 (M+H)⁺

Preparation 88

Methanesulfonyl chloride (1.4 ml) was added dropwise to a solution of acetic acid 3-amino-4-benzylphenyl ester (4.3 g) in pyridine (20 ml) under ice-cooling over 10 minutes, and the mixture was stirred at room temperature for a further 1 hour. Therein was added water (100 ml) and the resulting

mixture was stirred at the same temperature for 1 hour. The precipitates were collected by filtration, dissolved into chloroform (100 ml), followed by its dryness over magnesium sulfate and its evaporation in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford acetic acid 4-benzyloxy-3-methanesulfonylaminophenyl ester (1.6 g).

NMR (CDCl_3 , δ) : 2.27 (3H, s), 2.95 (3H, s), 5.09 (2H, s), 6.80-7.03 (3H, m), 7.25-7.45 (6H, m)

(+) APCI-MASS (m/z) : 336 (M+H)⁺

Preparation 89

A solution of acetic acid 4-benzyloxy-3-methanesulfonylaminophenyl ester (1.6 g) and potassium hydride (2.67 g) in methanol (10 ml) was stirred for 18 hours at room temperature. The reaction mixture was acidified with 1N hydrochloric acid to pH 5-7 and extracted with ethyl acetate. The extract was washed with brine, dried over magnesium sulfate, and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford N-(2-benzyloxy-5-hydroxyphenyl)methanesulfonamide (750 mg).

NMR (CDCl_3 , δ) : 2.90 (3H, s), 5.04 (2H, s), 6.58 (1H, dd, J=2.9 and 8.8Hz), 6.80-6.90 (2H, m), 7.09 (1H, d, J=2.9Hz), 7.30-7.50 (6H, m)

(+) APCI-MASS (m/z) : 294 (M+H)⁺

Preparation 90

Under nitrogen, to a solution of N-(2-benzyloxy-5-hydroxyphenyl)methanesulfonamide (740 mg) and sodium hydride (92.4 mg) in N,N-dimethylformamide (30 ml) was added (2S)-(+)-1-tosyloxy-2,3-epoxypropane (616 mg) at 0°C and the mixture was stirred at the same temperature for 0.5 hour. The mixture was allowed to warm to room temperature and stirred for 2.5 hours at this temperature. The resulting

mixture was poured into 10% aqueous ammonium chloride solution and extracted with ethyl acetate. The extract was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford (2S)-1-(4-benzyloxy-3-methanesulfonylamino)phenoxy-2,3-epoxypropane (440 mg).

NMR (CDCl₃, δ) : 2.75 (1H, dd, J=2.7 and 4.9Hz), 2.84-2.95 (4H, m), 3.30-3.37 (1H, m), 3.90 (1H, dd, J=5.8 and 11.1Hz), 4.07-4.25 (1H, m), 5.05 (2H, s), 6.63-7.48 (9H, m)

(+) APCI-MASS (m/z) : 350 (M+H)⁺

Preparation 91

To a solution of 2-[8-[(2S)-2-hydroxy-3-phenoxypropylamino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid ethyl ester (3.29 g) and triethylamine (2.55 ml) in tetrahydrofuran (20 ml) was added di-tert-butyl dicarbonate (1.53 g) under ice-cooling, and the mixture was stirred at room temperature for 18 hours. The resulting mixture was poured into saturated aqueous sodium bicarbonate solution and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford 2-[8-[N-tert-butoxycarbonyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid ethyl ester (3.61 g).

NMR (CDCl₃, δ) : 1.30 (3H, t, J=9.1Hz), 1.49 (9H, s), 1.50-2.05 (4H, m), 1.60-1.80 (3H, m), 3.20-3.80 (4H, m), 3.90-4.20 (3H, m), 4.27 (2H, q, J=7.1Hz), 4.57 (1H, s), 6.60-6.70 (2H, s), 6.90-7.05 (4H, m), 7.10-7.38 (2H, m)

Preparation 92

To a solution of 2-[8-[N-tert-butoxycarbonyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid ethyl ester (3.60 g) in ethanol (20 ml) was added 1N sodium hydroxide (7 ml) under ice-cooling, and the mixture was stirred at room temperature for 2 hours. The resulting mixture was acidified with hydrochloric acid to pH 2 and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford 2-[8-[N-tert-butoxycarbonyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid (3.2 g).

NMR (CDCl₃, δ) : 1.49 (9H, s), 1.90-2.05 (4H, m), 2.60-2.80 (3H, m), 3.20-3.75 (4H, m), 3.80-4.30 (3H, m), 4.61 (2H, s), 6.60-6.75 (2H, m), 6.80-7.05 (4H, m), 7.20-7.40 (2H, m)

Preparation 93

Under nitrogen, to a solution of N-benzyl-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (300 mg) and 1-[N-phenyl-N-benzyloxycarbonylamino]-2,3-epoxypropane (300 mg) in dichloromethane (5 ml) was added ytterbium(III) trifluoromethanesulfonate (66 mg) at room temperature, and the mixture was stirred at the same temperature for 24 hours. The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 10:1 to 4:1) to give N-[3-[N'-benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-2-hydroxypropyl]-N-benzyloxycarbonylphenylamine (307 mg).

(+) APCI-MASS (m/z) : 565 (M+H)⁺

Example 1

Under nitrogen, to a solution of N-benzyl-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (487 mg) and (2S)-3-phenoxy-1,2-epoxypropane (338 mg) in dichloromethane (10 ml) was added ytterbium(III) trifluoromethanesulfonate (107 mg) at room temperature, and the mixture was stirred at the same temperature for 3 hours. The resulting mixture was poured into saturated aqueous sodium bicarbonate solution, and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate, and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford (2S)-1-[N-benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol (750 mg).

NMR (CHCl_3 , δ) : 1.22-2.15 (4H, m), 2.45-3.10 (6H, m), 3.32-3.39 (1H, m), 3.70-4.10 (7H, m), 4.20-4.25 (1H, m), 6.58-7.00 (7H, m), 7.20-7.45 (6H, m)

MASS (m/z) : 432 (M+H)⁺

The following compounds [Example 2 to 4] were obtained according to a similar manner to that of Example 1.

Example 2

(2S)-1-[N-Benzyl-N-(1-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol

NMR (CHCl_3 , δ) : 1.10-2.35 (5H, m), 2.50-3.15 (6H, m), 3.32-3.48 (1H, m), 3.64-3.98 (5H, m), 3.77 (3H, s), 6.69-7.09 (6H, m), 7.18-7.32 (7H, m)

MASS (m/z) : 432 (M+H)⁺

Example 3

2-[8-[N'-Benzyl-(2-hydroxy-3-phenoxypropyl)amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-phenylacetamide

NMR (CDCl₃, δ) : 1.22-2.33 (4H, m), 2.53-3.18 (7H, m),
 3.67-4.05 (5H, m), 4.55-4.63 (2H, m), 6.65-6.78
 (1H, m), 6.78-7.05 (5H, m), 7.05-7.47 (10H, m),
 7.60 (2H, d, J=8.5Hz)

MASS (m/z) : 551 (M+H)⁺

Example 4

2-[8-[N'-Benzyl-(2-hydroxy-3-phenoxypropyl)amino]-
 6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-
 butylacetamide

NMR (CDCl₃, δ) : 0.92 (3H, t, J=7.4Hz), 1.15-2.30 (8H,
 m), 2.50-2.95 (6H, m), 2.98-3.15 (1H, m), 3.35 (2H,
 q, J=6.7Hz), 3.63-4.01 (5H, m), 4.41-4.50 (2H, m),
 6.47-7.03 (7H, m), 7.15-7.45 (7H, m)

MASS (m/z) : 531 (M+H)⁺

Example 5

N-Benzyl-(2-methoxy-6,7,8,9-tetrahydro-5H-
 benzocyclohepten-6-yl)amine hydrochloride was converted to
 the corresponding free base in a usual manner. A mixture of
 (2S)-3-phenoxy-1,2-epoxypropane (75 mg), N-benzyl-(2-methoxy-
 6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (140 mg),
 and ytterbium(III) trifluoromethanesulfonate (93 mg) in
 toluene (2.6 ml) was stirred at room temperature for 3 days
 and partitioned between ethyl acetate and aqueous sodium
 bicarbonate. The organic layer was separated, washed with
 brine, dried over sodium sulfate, and evaporated in vacuo.
 The residue was chromatographed (dichloromethane-methanol)
 over silica gel to afford (2S)-1-[N-benzyl-N-(2-methoxy-
 6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-
 2-propanol (200 mg).

IR (KBr) : 3427, 1248 cm⁻¹

NMR (CDCl₃, δ) : 1.25-2.25 (4H, m), 2.5-3.02 (7H, m),
 3.65-4.0 (8H, m), 6.63 (2H, m), 6.83-6.98 (3H, m),
 7.03-7.08 (1H, m), 7.18-7.31 (7H, m)

MASS (m/z) : 432 (M+H)⁺

Example 6

A mixture of (2S)-1-[N-benzyl-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol (750 mg) and 10% palladium on activated carbon (50% wet, 200 mg) in methanol (10 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 3 hours, and filtered. The filtrate was evaporated in vacuo and treated with 4N hydrogencchloride in ethyl acetate to afford (2S)-1-[(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol hydrochloride (446 mg).

NMR (DMSO-d₆, δ) : 1.09-1.30 (1H, m), 1.80-2.10 (2H, m), 2.25-2.34 (1H, m), 2.55-2.70 (2H, m), 3.00-3.70 (5H, m), 3.73 (3H, s), 4.01 (2H, d, J=4.0Hz), 4.20-4.30 (1H, m), 5.94 (1H, br s), 6.71 (1H, dd, J=2.6 and 8.2Hz), 6.83 (1H, dd, J=2.7 and 6.2Hz), 6.87-7.08 (4H, m), 7.23-7.36 (2H, m), 8.90 (1H, br s), 8.29 (1H, br s)

MASS (m/z) : 342 (M+H)⁺

The following compounds [Example 7 and 8] were obtained according to a similar manner to that of Example 6.

Example 7

(2S)-1-[(2-Methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol hydrochloride

IR (KBr) : 3423, 2800-2350, 1246 cm⁻¹

NMR (DMSO-d₆, δ) : 1.35 (1H, m), 1.75-2.05 (2H, m), 2.25-2.35 (1H, m), 2.70 (2H, m), 2.95-3.35 (5H, m), 3.71 (3H, s), 4.01 (2H, m), 4.24 (1H, m), 5.91 (1H, d, J=4.8Hz), 6.66-6.73 (2H, m), 6.92-6.70 (3H, m), 7.12 (1H, dd, J=8.0 and 2.5Hz), 7.31 (2H, t, J=7.8Hz), 8.78 (1H, br), 9.07 (1H, br)

MASS (m/z) : 342 (M+H)⁺

Example 8

(2S)-1-[(1-Methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol hydrochloride

NMR (DMSO-d₆, δ) : 1.04-1.32 (1H, m), 1.75-2.11 (2H, m), 2.23-2.40 (1H, m), 2.96-3.49 (7H, m), 3.75 (3H, s), 3.96-4.03 (2H, m), 4.26 (1H, br s), 5.92 (1H, br s), 6.81-7.01 (5H, m), 7.11 (1H, t, J=7.7Hz), 7.31 (2H, t, J=7.7Hz), 8.88 (1H, br s), 9.25 (1H, br s)

MASS (m/z) : 342 (M+H)⁺

Example 9

A mixture of 2-[8-[N'-benzyl-(2-hydroxy-3-phenoxypropyl)amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-phenylacetamide (360 mg) and 10% palladium on activated carbon (50% wet, 70 mg) in ethanol (5 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 4 hours and filtered. The filtrate was evaporated in vacuo. The residue was chromatographed (chloroform-methanol) over silica gel to afford 2-[8-(2-hydroxy-3-phenoxypropylamino)-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-phenylacetamide (306 mg).

The above obtained compound (306 mg) was treated with methanesulfonic acid (43 μl) to afford 2-[8-(2-hydroxy-3-phenoxypropylamino)-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-phenylacetamide methanesulfonate (303 mg).

NMR (DMSO-d₆, δ) : 1.18-1.43 (1H, m), 1.74-2.13 (2H, m), 2.18-2.37 (1H, m), 2.39 (3H, s), 2.60-2.83 (2H, m), 2.95-3.38 (5H, m), 3.95-4.40 (3H, m), 4.68 (2H, s), 6.78 (1H, d, J=8.0Hz), 6.86-7.14 (6H, m), 7.32 (4H, t, J=7.6Hz), 7.65 (2H, d, J=8.0Hz), 8.59 (2H, br s), 10.11 (1H, s)

MASS (m/z) : 461 (M+H)⁺

Example 10

A mixture of 2-[8-[N'-benzyl-(2-hydroxy-3-phenoxypropyl)amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-butylacetamide (206 mg) and 10% palladium on activated carbon (50% wet, 40 mg) in ethanol (5 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 4 hours, and filtered. The filtrate was evaporated in vacuo. The residue was chromatographed (chloroform-methanol) over silica gel to afford 2-[8-(2-hydroxy-3-phenoxypropylamino)-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-butylacetamide (171 mg).

The above obtained compound (70 mg) was treated with oxalic acid (14 mg) to afford 2-[8-(2-hydroxy-3-phenoxypropylamino)-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-butylacetamide oxalate (28 mg).

NMR (DMSO-d₆, δ) : 0.85 (3H, t, J=6.9Hz), 1.10-1.50 (5H, m), 1.70-2.12 (2H, m), 2.15-2.37 (1H, m), 2.60-2.78 (2H, m), 2.90-3.37 (7H, m), 3.99 (2H, br s), 4.20 (1H, br s), 4.63 (2H, s), 6.72 (1H, d, J=8.0Hz), 6.77-7.10 (5H, m), 7.31 (2H, d, J=7.4Hz)

MASS (m/z) : 441 (M+H)⁺

Example 11

A mixture of 4-oxiranylmethoxy-1,3-dihydro-2-benzimidazolone (82.5 mg) and N-benzyl-(6,7,8,9-tetrahydro-2-hydroxy-5H-benzocyclohepten-6-yl)amine (107 mg) in ethanol (1.5 ml) was stirred under reflux for 13 hours, cooled to room temperature and evaporated in vacuo. The residue was chromatographed (dichloromethane-methanol) by silica gel (4.2 g) to afford 1-[N-benzyl-N-(6,7,8,9-tetrahydro-2-hydroxy-5H-benzocyclohepten-6-yl)amino]-3-(1,3-dihydro-2-benzoimidazolone-4-yloxy)-2-propanol (123 mg) as a colorless powder.

mp : 111-133°C (dec.)

IR (KBr) : 3408 (br), 3257 (br), 1695, 1248 cm⁻¹

NMR (DMSO- d_6 , δ) : 1.03-1.97 (4H, m), 2.22-2.84 (6H, m), 3.63-4.17 (6H, m), 4.60 and 4.67 (1H, each d, $J=4.1\text{Hz}$), 6.43 (2H, m), 6.53-6.63 (2H, m), 6.79-6.95 (2H, m), 7.11-7.35 (5H, m), 8.99 (1H, s),
10.55 (2H, br s)
(+) APCI-MASS (m/z) : 474 (M+H)⁺

Example 12

A solution of (2S)-1,2-epoxy-3-phenoxypropane (53 mg) (IL FARMACO, 50 (10), 643 (1995)) and N-benzyl-(3-chloro-6,7,8,9-tetrahydro-2-hydroxy-5H-benzocyclohepten-6-yl)amine (89 mg) in ethanol (1 ml) was refluxed for 19 hours, cooled to room temperature and evaporated in vacuo. The residue was chromatographed (dichloromethane-methanol) by silica gel (2.9 g), and the eluate was treated with 4N hydrogen chloride in ethyl acetate to afford (2S)-1-[N-benzyl-N-(3-chloro-6,7,8,9-tetrahydro-2-hydroxy-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol hydrochloride (149 mg) as a colorless powder.

mp : 94-99°C (dec.)

IR (KBr) : 3421 (br), 2750-2450 (m), 1244 cm^{-1}

Example 13

Under nitrogen, to a solution of N-benzyl-(3-isopropoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (258 mg) and (2S)-3-phenoxy-1,2-epoxypropane (138 mg) in dichloromethane (5 ml) was added ytterbium(III) trifluoromethanesulfonate (52 mg) at room temperature, and the mixture was stirred at the same temperature for 60 hours. The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 10:1 to 5:1) to give (2S)-1-[N-benzyl-N-(3-

isopropoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-
amino]-3-phenoxy-2-propanol (303 mg).

(+) APCI-MASS (m/z) : 460 (M+H)⁺

- 5 The following compounds [Examples 14 to 27] were
obtained according to a similar manner to that of Example 13.

Example 14

10 (2S)-1-[N-Benzyl-N-(3-hexyloxy-6,7,8,9-tetrahydro-5H-
benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol

(+) APCI-MASS (m/z) : 502 (M+H)⁺

Example 15

15 (2S)-1-[N-Benzyl-N-(3-benzyloxy-6,7,8,9-tetrahydro-5H-
benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol

(+) ESI-MASS (m/z) : 508 (M+H)⁺

Example 16

20 (2S)-1-[N-Benzyl-N-[3-(3-cyanopyridin-2-yloxy)-6,7,8,9-
tetrahydro-5H-benzocyclohepten-6-yl]amino]-3-phenoxy-2-
propanol

(+) APCI-MASS (m/z) : 520 (M+H)⁺

Example 17

25 (2S)-1-[N-Benzyl-N-[3-(2-methoxyethoxy)-6,7,8,9-
tetrahydro-5H-benzocyclohepten-6-yl]amino]-3-phenoxy-2-
propanol

(+) APCI-MASS (m/z) : 476 (M+H)⁺

30 Example 18

(2S)-1-[N-Benzyl-N-(3-nitro-6,7,8,9-tetrahydro-5H-
benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol

(+) APCI-MASS (m/z) : 447 (M+H)⁺

35 Example 19

(2S)-1-[N-Benzyl-N-(6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol

(+) APCI-MASS (m/z) : 402 (M+H)⁺

5 Example 20

(2S)-1-[N-Benzyl-N-(3-phenyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol

(+) ESI-MASS (m/z) : 478 (M+H)⁺

10 Example 21

(2S)-1-[N-Benzyl-N-(2-chloro-3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol

(+) ESI-MASS (m/z) : 466, 468 (M+H)⁺

15

Example 22

(2S)-1-[N-Benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenylthio-2-propanol

(+) APCI-MASS (m/z) : 448 (M+H)⁺

20

Example 23

(2S)-1-[N-Benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-(4-fluorophenoxy)-2-propanol

NMR (CDCl₃, δ) : 0.85-1.25 (1H, m), 1.65-2.1 (2H, m),
2.2-2.4 (1H, m), 2.45-2.95 (7H, m), 3.65-3.85 (7H, m),
3.85-4.1 (1H, m), 6.55-6.6 (1H, m), 6.65-6.75 (1H, m), 6.8-7.4 (10H, m)

(+) APCI-MASS (m/z) : 450 (M+H)⁺

25

30 Example 24

(2S)-1-[N-Benzyl-N-(6,7,8,9-tetrahydro-4-methoxy-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol hydrochloride

NMR (DMSO-d₆, δ) : 1.10-1.38 (1H, m), 1.97-2.20 (2H, m),
2.38-2.52 (1H, m), 2.60-3.00 (4H, m), 3.18-3.58

35

(3H, m), 3.70-4.02 (3H, m), 3.79 (3H, s), 4.45-4.64
 (2H, m), 6.01 (1H, br), 6.70-6.99 (5H, m), 7.13
 (1H, dd, J=8.0, 8.0Hz), 7.30 (2H, dd, J=7.9,
 7.9Hz), 7.40-7.56 (3H, m), 7.65-7.80 (2H, m), 9.42-
 10.09 (1H, br)

(+) ESI-MASS (m/z) : 432 (M+H)⁺

Example 25

(2S)-1-[N-Benzyl-N-(3-methoxycarbonyl-6,7,8,9-
 tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-
 propanol hydrochloride

NMR (DMSO-d₆, δ) : 1.10-1.33 (1H, m), 1.99-2.21 (2H,
 m), 2.40-2.55 (1H, m), 2.75-2.91 (2H, m), 3.04-3.17
 (1H, m), 3.26-3.38 (3H, m), 3.45-3.59 (2H, m),
 3.70-4.01 (2H, m), 3.84 (3H, s), 4.39-4.69 (2H, m),
 5.92-6.05 (1H, br), 6.81-7.00 (3H, m), 7.23-7.34
 (3H, m), 7.38-7.48 (3H, m), 7.68-7.97 (4H, m),
 9.67-9.95 (1H, br)

(+) ESI-MASS (m/z) : 460 (M+H)⁺

Example 26

(2S)-1-[N-Benzyl-N-(3-carbamoyl-6,7,8,9-tetrahydro-5H-
 benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol
 hydrochloride

NMR (DMSO-d₆, δ) : 1.18-1.34 (1H, m), 2.00-2.20 (2H,
 m), 2.33-2.58 (1H, m), 2.78-2.90 (2H, m), 3.02-3.50
 (5H, m), 3.72-4.02 (3H, m), 4.28-4.62 (2H, m), 5.98
 (1H, br), 6.80-6.98 (3H, m), 7.17-7.44 (7H, m),
 7.63-7.91 (5H, m), 9.36-9.92 (1H, br)

(+) APCI-MASS (m/z) : 445 (M+H)⁺

Example 27

1-[N-Benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-
 benzocyclohepten-6-yl)amino]-4-phenyl-2-butanol

(+) APCI-MASS (m/z) : 430 (M+H)⁺

Example 28

Under nitrogen, to a solution of N-(3-fluoromethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-tert-butoxycarbonyl-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-amine (300 mg) in tetrahydrofuran (5 ml) was added tetra-n-butylammonium fluoride (1M in tetrahydrofuran, 0.58 ml) at 5°C, and the mixture was stirred at the same temperature for 15 minutes. The resulting mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 3:1) to give N-(3-fluoromethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-tert-butoxycarbonyl-[(2S)-2-hydroxy-3-phenoxypropyl]amine (227 mg).

NMR (CDCl₃, δ) : 1.1-1.6 (1H, m), 1.49 (9H, s), 1.9-2.1 (3H, m), 2.65-2.75 (3H, m), 3.15-3.7 (4H, m), 3.8-4.2 (3H, m), 5.67 (2H, d, J=54.8Hz), 6.75-7.1 (5H, m), 7.25-7.3 (2H, m)

Example 29

Under nitrogen, to a suspension of sodium hydride (60% in oil, 12 mg) in N,N-dimethylformamide (3 ml) was added N-(3-hydroxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-tert-butoxycarbonyl-[(2S)-3-phenoxy-2-(triethylsilyloxy)-propyl]amine (150 mg) at 5°C. After being stirred at the same temperature for 1.5 hours, to this one was added dimethylsulfonyl chloride (31 μl), and the mixture was stirred at 60°C for 15 hours. The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 3:1) to give (1): N-(3-dimethylsulfonyloxy-6,7,8,9-tetrahydro-5H-

benzocyclohepten-6-yl)-N-tert-butoxycarbonyl-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]amine (59 mg) and (2): N-(3-dimethylsulfonyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-tert-butoxycarbonyl-[(2S)-2-hydroxy-3-phenoxypropyl]-amine (66 mg).

(1): NMR (CDCl₃, δ) : 0.55-0.7 (6H, m), 0.9-1.05 (9H, m), 0.2-1.5 (1H, m), 1.48 (9H, m), 1.9-2.15 (3H, m), 2.7-3.0 (3H, m), 2.95 (6H, m), 3.1-3.7 (4H, m), 3.85-4.05 (2H, m), 4.3-4.4 (1H, m), 6.85-7.1 (6H, m), 7.25-7.3 (2H, m)

(+) APCI-MASS (m/z) : 549 (M-Boc+2H)⁺

(2): NMR (CDCl₃, δ) : 0.55-0.7 (6H, m), 0.9-1.05 (9H, m), 1.2-1.5 (1H, m), 1.48 (9H, m), 1.9-2.15 (3H, m), 2.7-3.0 (3H, m), 2.95 (6H, m), 3.1-3.7 (4H, m), 3.85-4.05 (2H, m), 4.3-4.4 (1H, m), 6.85-7.1 (6H, m), 7.25-7.3 (2H, m)

(+) APCI-MASS (m/z) : 435 (M-Boc+2H)⁺

Example 30

Under nitrogen, to a solution of 2-(8-benzylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy)acetamide (260 mg) and (2S)-3-phenoxy-1,2-epoxypropane (130 mg) in dichloromethane (3 ml) was added ytterbium(III) trifluoromethanesulfonate (50 mg) at room temperature, and the mixture was stirred at the same temperature for 12 hours. The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate and extracted with dichloromethane. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (chloroform:ethyl acetate = 20:1 to 10:1), followed by treatment with methanesulfonic acid in chloroform to give 2-[8-[N-benzyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetamide

methanesulfonate (290 mg).

(+) APCI-MASS (m/z) : 475 (M-MsOH+H)⁺

Example 31

5 A solution of 2-[8-[N-benzyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid ethyl ester (200 mg) and N,N-dimethylethylenediamine (130 μ l) in methanol (2 ml) was stirred at room temperature for 24 hours. After evaporation
10 in vacuo, the residue was purified by preparative thin layer chromatography (silica gel, chloroform:methanol = 10:1) to give 2-[8-[N'-benzyl-N'-[(2S)-2-hydroxy-3-phenoxypropyl]-amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-(2-dimethylaminoethyl)acetamide (215 mg).

15 (+) APCI-MASS (m/z) : 546 (M+H)⁺

Example 32

Under nitrogen, a solution of 2-[8-[N-tert-butoxycarbonyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-
20 6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid (100 mg) in dichloromethane (3 ml) were added N-methylbutylamine (29 μ l) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (47 mg) in the presence of catalytic amounts of 4-dimethylaminopyridine at 5°C, and the
25 mixture was stirred at room temperature for 12 hours. The resulting mixture was poured into 1N hydrochloric acid and extracted with ethyl acetate. The organic layer was successively washed with saturated aqueous sodium hydrogencarbonate and brine, dried over magnesium sulfate and
30 evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 1:1) to give N-[3-[(butylmethylcarbamoyl)methoxy]-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]-N-tert-butoxycarbonyl-[(2S)-2-hydroxy-3-phenoxypropyl]amine (96 mg).

35 NMR (CHCl₃, δ) : 0.92 (3H, q, J=7.6Hz), 1.18-1.70 (5H,

m), 1.49 (9H, s), 1.93-2.15 (3H, m), 2.59-2.82 (3H, m), 2.94-3.04 (3H, m), 3.20-3.66 (6H, m), 3.83-4.25 (3H, m), 4.62-4.67 (2H, m), 6.65-6.77 (2H, m), 6.87-7.02 (4H, m), 7.25-7.34 (2H, m)

5 (+) APCI-MASS (m/z) : 455 (M-Boc+2H)⁺

The following compounds[Examples 33 to 38] were obtained according to a similar manner to that of Example 32.

10 Example 33

N-[(2S)-2-Hydroxy-3-phenoxypropyl]-N-tert-butoxycarbonyl-[3-(sec-butylcarbamoylmethoxy)-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]amine

(+) APCI-MASS (m/z) : 441 (M-Boc+2H)⁺

15

Example 34

N-(3-Cyclohexylcarbamoylmethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-tert-butoxycarbonyl-[(2S)-2-hydroxy-3-phenoxypropyl]amine

20 (+) APCI-MASS (m/z) : 467 (M-Boc+2H)⁺

Example 35

N-[3-(Benzylcarbamoylmethoxy)-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]-N-tert-butoxycarbonyl-[(2S)-2-hydroxy-3-phenoxypropyl]amine

25

(+) APCI-MASS (m/z) : 475 (M-Boc+2H)⁺

Example 36

N-[(2S)-2-Hydroxy-3-phenoxypropyl]-N-tert-butoxycarbonyl-[3-[(2-methylthioethylcarbamoyl)methoxy]-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]amine

30

(+) APCI-MASS (m/z) : 459 (M-Boc+2H)⁺

Example 37

35 N-[(2S)-2-Hydroxy-3-phenoxypropyl]-N-tert-

butoxycarbonyl-[3-(2-piperidinocarbonylethoxy)-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]amine

(+) APCI-MASS (m/z) : 453 (M-Boc+2H)⁺

5 Example 38

N-[(2S)-2-Hydroxy-3-phenoxypropyl]-N-tert-butoxycarbonyl-[3-[(1H-indol-5-yl-carbamoyl)methoxy]-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]amine

10 NMR (CDCl₃, δ) : 1.22-1.70 (1H, m), 1.50 (9H, s), 1.96-2.18 (3H, m), 2.66-2.88 (3H, m), 3.26-3.70 (4H, m), 3.85-4.26 (3H, m), 4.59 (2H, s), 6.51-6.56 (1H, m), 6.68-6.87 (2H, m), 6.88-7.09 (4H, m), 7.23-7.39 (6H, m), 7.92 (1H, m), 8.16-8.37 (2H, m)

(+) APCI-MASS (m/z) : 500 (M-Boc+2H)⁺

15

Example 39

A suspension of powdered iron (160 mg) and ammonium chloride (15 mg) in a mixture of ethanol (6 ml) and water (2 ml) was refluxed, and to this one was dropwise added (2S)-1-[N-benzyl-N-(3-nitro-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol (158 mg) in ethanol (3 ml). After being further refluxed for 1 hour, insoluble materials were filtered off. The filtrate was evaporated in vacuo. The residue was dissolved into a mixture of saturated aqueous sodium hydrogencarbonate and ethyl acetate. After separation, the organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 2:1 to 1:1) to give (2S)-1-[N-(3-amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-benzylamino]-3-phenoxy-2-propanol (122 mg).

30

(+) ESI-MASS (m/z) : 417 (M+H)⁺

Example 40

35

Under nitrogen, a solution of (2S)-1-[N-(3-amino-6,7,8,

9-tetrahydro-5H-benzocyclohepten-6-yl)-N-benzylamino]-3-phenoxy-2-propanol (200 mg), methyl iodide (0.15 ml) and N,N-diisopropylethylamine (0.25 ml) in dichloromethane (5 ml) was stirred at room temperature for 12 hours. The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 4:1) to give a less polar diastereomer: (2S)-1-[N-benzyl-N-(3-dimethylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol (28 mg) and a more polar diastereomer: (2S)-1-[N-benzyl-N-(3-dimethylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol (25 mg).

a less polar diastereomer

(+) APCI-MASS (m/z) : 445 (M+H)⁺

a more polar diastereomer

(+) APCI-MASS (m/z) : 445 (M+H)⁺

Example 41

Under nitrogen, a solution of (2S)-1-[N-(3-amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-benzylamino]-3-phenoxy-2-propanol (113 mg), acetic anhydride (28 µl) and pyridine (24 µl) in dichloromethane (5 ml) was stirred at 5°C for 1 hour. The resulting mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 1:1 to 1:2) to give N-benzyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]-3-acetyl-amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (92 mg).

(+) APCI-MASS (m/z) : 459 (M+H)⁺

Example 42

Under nitrogen, a solution of (2S)-1-[N-(3-amino-
 6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-benzylamino]-
 3-phenoxy-2-propanol (150 mg), benzoyl chloride (44 μ l),
 pyridine (35 μ l), and catalytic amounts of
 4-dimethylaminopyridine in dichloromethane (5 ml) was stirred
 at room temperature for 5 hours. The resulting mixture was
 poured into saturated aqueous sodium hydrogencarbonate and
 extracted with ethyl acetate. The organic layer was washed
 with brine, dried over anhydrous magnesium sulfate and
 evaporated in vacuo. The residue was purified by column
 chromatography on silica gel (hexane:ethyl acetate = 2:1) to
 give N-benzyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]-3-
 benzoylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylamine
 (163 mg).

(+) APCI-MASS (m/z) : 521 (M+H)⁺

Example 43

Under nitrogen, a solution of (2S)-1-[N-(3-amino-
 6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-benzylamino]-
 3-phenoxy-2-propanol (150 mg), methanesulfonyl chloride (29
 μ l), pyridine (35 μ l), and catalytic amounts of 4-
 dimethylaminopyridine in dichloromethane (5 ml) was stirred
 at room temperature for 4 hours. The resulting mixture was
 poured into saturated aqueous sodium hydrogencarbonate and
 extracted with ethyl acetate. The organic layer was washed
 with brine, dried over anhydrous magnesium sulfate and
 evaporated in vacuo. The residue was purified by column
 chromatography on silica gel (hexane:ethyl acetate = 2:1) to
 give N-benzyl-N-[(2S)-2-hydroxy-3-phenoxypropylamino]-3-
 methylsulfonylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-
 ylamine (178 mg).

(+) APCI-MASS (m/z) : 495 (M+H)⁺

The following compound was obtained according to a similar manner to that of Example 43.

Example 44

5 N-Benzyl-N-[(2S)-2-hydroxy-3-phenoxypropylamino]-
6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylamine
 (+) ESI-MASS (m/z) : 557 (M+H)⁺

Example 45

10 Under nitrogen, a solution of (2S)-1-[N-(3-amino-
6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-benzylamino]-
3-phenoxy-2-propanol (150 mg), methyl chloroformate (29 μ l),
pyridine (25 μ l) and catalytic amounts of
4-dimethylaminopyridine in dichloromethane (3 ml) was stirred
15 at room temperature for 1.5 hours. The resulting mixture was
poured into saturated aqueous sodium hydrogencarbonate and
extracted with ethyl acetate. The organic layer was washed
with brine, dried over anhydrous magnesium sulfate and
evaporated in vacuo. The residue was purified by column
20 chromatography on silica gel (hexane:ethyl acetate = 3:1) to
give (2S)-1-[N-(3-methoxycarbonylamino-6,7,8,9-tetrahydro-5H-
benzocyclohepten-6-yl)-N-benzylamino]-3-phenoxy-2-propanol
(158 mg).
 (+) APCI-MASS (m/z) : 475 (M+H)⁺

25

The following compound was obtained according to a similar manner to that of Example 46.

Example 46

30 (2S)-1-[N-(3-phenoxy-carbonylamino-6,7,8,9-tetrahydro-5H-
benzocyclohepten-6-yl)-N-benzylamino]-3-phenoxy-2-propanol
 (+) APCI-MASS (m/z) : 537 (M+H)⁺

Example 47

35 Under nitrogen, a solution of (2S)-1-[N-(3-amino-

6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-benzylamino]-3-phenoxy-2-propanol (150 mg), dimethylaminosulfonyl chloride (43 μ l) and pyridine (31 μ l) in toluent (5 ml) was stirred at 80°C for 24 hours. The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 2:1) to give (2S)-1-[N-(3-dimethylaminosulfonylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-benzylamino]-3-phenoxy-2-propanol (156 mg).

(+) APCI-MASS (m/z) : 524 (M+H)⁺

The following compound was obtained according to a similar manner to that of Example 47.

Example 48

(2S)-1-[N-(3-Sulfamoylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-benzylamino]-3-phenoxy-2-propanol

(+) APCI-MASS (m/z) : 496 (M+H)⁺

Example 49

To a solution of (2S)-1-[N-benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenylthio-2-propanol (221 mg) in methanol (10 ml) was added OXONE (610 mg) in water (3 ml) at room temperature, and the mixture was stirred at the same temperature for 5 hours. The resulting mixture was filtered off to remove precipitates and the filtrate was evaporated in vacuo. The residue was dissolved into a mixture of saturated aqueous sodium hydrogencarbonate and ethyl acetate. After separation, the organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (hexane:ethyl acetate = 2:1) to

give (2S)-1-[N-benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenylsulfonyl-2-propanol (170 mg).

(+) APCI-MASS (m/z) : 480 (M+H)⁺

5

Example 50

A mixture of N-benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (200 mg) and 10% palladium on activated carbon (50% wet, 50 mg) in methanol (5 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 3 hours. After filtration, the filtrate was evaporated in vacuo. A mixture of the residue and [5-[(2S)-oxiranyl)methoxypyridin-2-yl]carbamic acid benzyl ester (174 mg) in methanol (10 ml) was refluxed for 20 hours. After removal of the solvent in vacuo, the residue was purified by column chromatography on silica gel (chloroform:methanol = 50:1 to 20:1) to give N-[5-[(2S)-2-hydroxy-3-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-ylamino)propoxy]pyridin-2-yl]benzyloxycarbonylamine (200 mg).

(+) APCI-MASS (m/z) : 492 (M+H)⁺

The following compound was obtained according to a similar manner to that of Example 50.

25

Example 51

4-[2-Hydroxy-3-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-ylamino)propoxy]-1,3-dihydrobenzimidazol-2-one hydrochloride

(+) APCI-MASS (m/z) : 398 (M-HCl+H)⁺

30

Example 52

8-[[[(2S)-2-Hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxylic acid methyl ester hydrochloride (8.40 g) was converted to the

35

corresponding free base in a usual manner. The free base was dissolved in tetrahydrofuran (60 ml), and a solution of di-tert-butyl dicarbonate (4.52 g) in tetrahydrofuran (10 ml) was added to the mixture. After stirring overnight, the reaction mixture was evaporated in vacuo, and the residue was purified by column chromatography on silica gel eluting with a mixture of hexane and ethyl acetate (5:1 to 3:1) to afford 8-[N-tert-butoxycarbonyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]-amino]-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxylic acid methyl ester (9.88 g).

NMR (DMSO- d_6 , δ) : 1.15-1.40 (1H, m), 1.41 (9H, s), 1.86-2.35 (3H, m), 2.68-3.60 (7H, m), 3.83 (3H, s), 3.85-4.08 (3H, m), 5.19 (1H, d, $J=5.1\text{Hz}$), 6.89-6.99 (3H, m), 7.22-7.33 (3H, m), 7.70 (1H, dd, $J=7.7$ and 1.6Hz), 7.75 (1H, s)

(+) ESI-MASS (m/z) : 492 (M+Na)⁺

Example 53

To a solution of 8-[N-tert-butoxycarbonyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxylic acid methyl ester (4.5 g) in methanol (45 ml) and tetrahydrofuran (15 ml) was added 1N sodium hydroxide aqueous solution. After stirring at room temperature for 6 hours, the reaction mixture was concentrated under reduced pressure, neutralized with 1N hydrochloric acid solution under ice-cooling. The product was extracted with ethyl acetate, and the organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo to afford 8-[N-tert-butoxycarbonyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxylic acid (4.48 g).

NMR (DMSO- d_6 , δ) : 1.19-1.40 (1H, m), 1.41 (9H, s), 1.89-2.37 (3H, m), 2.71-3.58 (7H, m), 3.80-4.03 (3H, m), 5.17 (1H, d, $J=5.0\text{Hz}$), 6.88-6.95 (3H, m), 7.19-7.32 (3H, m), 7.67 (1H, d, $J=7.7\text{Hz}$), 7.74 (1H,

s)

(+) APCI-MASS (m/z) : 356 (M-Boc+2H)⁺Example 54

To a mixture of 8-[N-tert-butoxycarbonyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxylic acid (0.30 g), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.15 g) and 1-hydroxybenzotriazole (0.11 g) in dichloromethane (6 ml) was added a solution of methylamine in tetrahydrofuran (2.0M, 0.40 ml). The reaction mixture was stirred at room temperature overnight, and partitioned between ethyl acetate and aqueous sodium bicarbonate solution. The organic layer was separated, washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel eluting with a mixture of chloroform and methanol (100:1 to 10:1) to afford N-[(2S)-2-hydroxy-3-phenoxypropyl]-N-tert-butoxycarbonyl-(3-methylcarbamoyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (0.13 g).

NMR (DMSO-d₆, δ) : 1.16-1.35 (1H, m), 1.40 (9H, s), 1.87-2.24 (3H, m), 2.76 (3H, d, J=4.6Hz), 2.70-2.92 (2H, m), 3.03-3.56 (5H, m), 3.85-4.02 (3H, m), 5.14 (1H, d, J=5.1Hz), 6.89-6.95 (3H, m), 7.15 (1H, d, J=7.7Hz), 7.28 (2H, dd, J=7.7 and 7.7Hz), 7.55 (1H, d, J=7.7Hz), 7.63 (1H, s), 8.28 (1H, t-like, J=4.6Hz)

(+) ESI-MASS (m/z) : 491 (M+Na)⁺

The following compounds [Examples 55 and 56] were obtained according to a similar manner to that of Example 54.

Example 55

N-[(2S)-2-Hydroxy-3-phenoxypropyl]-N-tert-butoxycarbonyl-(3-dimethylcarbamoyl-6,7,8,9-tetrahydro-5H-

benzocyclohepten-6-yl)amine

NMR (DMSO- d_6 , δ) : 1.19-1.38 (1H, m), 1.40 (9H, s),
 1.90-2.08 (2H, m), 2.12-2.30 (1H, m), 2.68-3.10
 (4H, m), 2.93 (6H, s), 3.18-3.60 (3H, m), 3.85-4.08
 (3H, m), 5.16 (1H, br), 6.88-6.95 (3H, m), 7.12-
 7.17 (3H, m), 7.28 (2H, dd, $J=8.0$ and 8.0Hz)
 (+) ESI-MASS (m/z) : 505 (M+Na)⁺

Example 56

N-[(2S)-2-Hydroxy-3-phenoxypropyl]-N-tert-
 butoxycarbonyl-(3-phenylcarbamoyl-6,7,8,9-tetrahydro-5H-
 benzocyclohepten-6-yl)amine

NMR (DMSO- d_6 , δ) : 1.19-1.20 (1H, m), 1.41 (9H, s),
 1.88-2.36 (3H, m), 2.75-3.60 (7H, m), 3.85-4.10
 (3H, m), 5.19 (1H, d, $J=5.0\text{Hz}$), 6.85-6.93 (3H, m),
 7.09 (1H, dd, $J=7.4$ and 7.4Hz), 7.22-7.39 (5H, m),
 7.64-7.78 (4H, m), 10.11 (1H, s)
 (+) ESI-MASS (m/z) : 553 (M+Na)⁺

Example 57

To a solution of 8-[N-benzyl-N-[(2S)-2-hydroxy-3-
 phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocycloheptene-
 2-carboxamide (3.00 g) in dry tetrahydrofuran (90 ml) was
 gradually added lithium aluminum hydride (1.02 g) under ice-
 cooling and nitrogen atmosphere. The mixture was then heated
 to reflux for 3 hours. Saturated sodium bicarbonate solution
 was added dropwise to the reaction mixture under ice-cooling,
 and the resulting precipitate was filtered off and washed
 with ethyl acetate. The filtrate was extracted with ethyl
 acetate, and the organic layer was washed with water and
 brine, dried over magnesium sulfate and evaporated in vacuo.
 The residue was purified by column chromatography on silica
 gel eluting with a mixture of chloroform and methanol (30:1
 to 5:1) to afford (2S)-1-[N-(3-aminomethyl-6,7,8,9-
 tetrahydro-5H-benzocyclohepten-6-yl)-N-benzylamino]-3-

phenoxy-2-propanol (2.75 g).

NMR (DMSO- d_6 , δ) : 1.00-1.20 (1H, m), 1.67-2.10 (3H, m), 2.23-2.40 (1H, m), 2.50-3.10 (8H, m), 3.62 (2H, d, $J=3.0\text{Hz}$), 3.70-3.86 (4H, m), 3.92-4.02 (1H, m), 4.78 (1H, br), 6.83-7.00 (5H, m), 7.05-7.38 (8H, m)
 (+) APCI-MASS (m/z) : 431 (M+H)⁺

Example 58

To a mixture of (2S)-1-[N-(3-aminomethyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-benzylamino]-3-phenoxy-2-propanol (0.30 g) and triethylamine (0.107 ml) in dichloromethane (6 ml) was added dropwise acetyl chloride (0.05 ml) under ice-cooling. The mixture was stirred at room temperature overnight and partitioned between water and ethyl acetate. The organic layer was separated, washed with water and brine, dried over magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel eluting with a mixture of chloroform and methanol (50:1). The product was treated with 4N hydrogen chloride in ethyl acetate and powdered from diisopropyl ether to afford N-[6-[N'-benzyl-N'-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-3-ylmethyl]acetamide (0.29 g).

NMR (DMSO- d_6 , δ) : 1.09-1.30 (1H, m), 1.86-1.90 (total 3H, s), 1.98-2.18 (2H, m), 2.38-2.62 (1H, m), 2.70-2.82 (2H, m), 3.03-3.68 (5H, m), 3.74-4.04 (3H, m), 4.10-4.24 (2H, m), 4.40-4.68 (2H, m), 6.01 (1H, br), 6.83-7.10 (6H, m), 7.24-7.34 (2H, m), 7.40-7.52 (3H, m), 7.65-7.82 (2H, m), 8.37 (1H, t, $J=6.0\text{Hz}$), 9.68-10.24 (1H, br)
 (+) APCI-MASS (m/z) : 473 (M+H)⁺

The following compounds [Example 59 to 61] were obtained according to a similar manner to that of Example 58.

Example 59

N-[8-[N'-Benzyl-N'-[(2S)-2-hydroxy-3-phenoxypropyl]-amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylmethyl]-benzamide hydrochloride

5 NMR (DMSO-d₆, δ) : 1.09-1.28 (1H, m), 1.93-2.19 (2H, m), 2.40-2.60 (1H, m), 2.67-2.84 (2H, m), 2.98-3.50 (5H, m), 3.74-4.02 (3H, m), 4.32-4.56 (4H, m), 6.00 (1H, br), 6.83-6.98 (3H, m), 7.07-7.53 (11H, m), 7.66-7.78 (2H, m), 7.94 (2H, dd, J=8.3 and 8.3Hz), 9.12 (1H, t, J=6.0Hz), 9.81-10.38 (1H, br)
10 (+) APCI-MASS (m/z) : 535 (M+H)⁺

Example 60

15 N-[8-[N'-Benzyl-[N'-(2S)-2-hydroxy-3-phenoxypropyl]-amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylmethyl]-methanesulfonamide hydrochloride

20 NMR (DMSO-d₆, δ) : 1.08-1.23 (1H, m), 1.97-2.18 (2H, m), 2.39-2.60 (1H, m), 2.66-2.80 (2H, m), 2.82, 2.89 (total 3H, s), 3.00-3.64 (5H, m), 3.74-4.13 (5H, m), 4.40-4.70 (2H, m), 5.95 (1H, br), 6.83-6.98 (3H, m), 7.09-7.60 (9H, m), 7.67-7.78 (2H, m), 9.73-10.36 (1H, br)
(+) APCI-MASS (m/z) : 509 (M+H)⁺

25 Example 61

N-[8-[N'-Benzyl-N'-[(2S)-2-hydroxy-3-phenoxypropyl]-amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylmethyl]-benzenesulfonamide hydrochloride

30 NMR (DMSO-d₆, δ) : 1.05-1.23 (1H, m), 1.96-2.18 (2H, m), 2.41-2.85 (3H, m), 2.99-3.48 (5H, m), 3.70-4.03 (5H, m), 4.45-4.70 (2H, m), 5.98-6.06 (1H, br), 6.83-7.06 (5H, m), 7.23-7.34 (3H, m), 7.40-7.84 (10H, m), 8.12-8.22 (1H, m), 9.71-10.31 (1H, br)
(+) APCI-MASS (m/z) : 571 (M+H)⁺

Example 62

A mixture of (2R)-1-(3-methylsulfonylamino-4-benzyloxy)phenyl-2,3-epoxypropane (39 mg), (RS)-N-benzyl-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (58 mg), ytterbium triflate (113 mg) and dichloromethane (1 ml) was stirred at room temperature overnight. The reaction mixture was washed up in the usual manner. The crude product was dissolved in isopropylalcohol (1.5 ml), heated with refluxing for 4.5 hours, evaporated and purified by column chromatography (silica gel, 20 cm³, eluent; 20% ethyl acetate-toluene) to give (2R)-N-[3-(3-methylsulfonylamino-4-benzyloxy)phenyl-2-hydroxypropyl]-N-[(6RS)-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)]benzylamine (40.2 mg) as an oil.

IR (Film) : 3272, 2927, 1610, 1508, 1456, 1332, 1261, 1161, 1122, 975 cm⁻¹

NMR (CDCl₃, δ) : 1.4-1.8 (4H, m), 2.4-3.1 (11H, m), 2.88 (3H, d, J=2.4Hz), 3.5-3.8 (1H, m), 3.79 (3H, s), 5.08 (2H, s), 6.6-6.8 (2H, m), 6.9-7.0 (3H, m), 7.1-7.5 (11H, m)

Example 63

Under nitrogen, to a solution of N-benzyl-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (120 mg) and (2S)-1-(4-acetylamino)phenoxy-2,3-epoxypropane (89 mg) in dichloromethane (10 ml) was added ytterbium(III) trifluoromethanesulfonate (50 mg) at room temperature, and the mixture was stirred at the same temperature for 3 hours. The resulting mixture was poured into saturated aqueous sodium bicarbonate solution and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford (2S)-1-[N-benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-(4-acetylaminophenoxy)-2-

propanol (80 mg).

(+) APCI-MASS (m/z) : 489 (M+H)⁺

Example 64

5 Under nitrogen, to a solution of 2-(8-benzylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy)-N-butylacetamide (200 mg) and (2S)-1-(1H-indol-5-yloxy)-2,3-epoxypropane (130 mg) in dichloromethane (10 ml) was added
10 ytterbium(III) trifluoromethanesulfonate (33 mg) at room temperature, and the mixture was stirred at the same temperature for 3 hours. The resulting mixture was poured into saturated aqueous sodium bicarbonate solution and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in
15 vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford 2-[8-[N-benzyl-N-[(2S)-2-hydroxy-3-(1H-indol-5-yloxy)propyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-butylacetamide (160 mg).

(+) APCI-MASS (m/z) : 570 (M+H)⁺

Example 65

under nitrogen, to a solution of 2-(8-benzylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy)-N-butylacetamide (388 mg) and (2S)-1-(4-benzyloxyphenoxy)-2,3-
25 epoxypropane (340 mg) in dichloromethane (10 ml) was added ytterbium(III) trifluoromethanesulfonate (63 mg) at room temperature, and the mixture was stirred at the same temperature for 3 hours. The resulting mixture was poured into saturated aqueous sodium bicarbonate solution and
30 extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford 2-[8-[N-benzyl-N-[(2S)-3-(4-benzyloxyphenoxy)-2-hydroxypropyl]amino]-6,7,8,9-
35 tetrahydro-5H-benzocyclohepten-2-yloxy]-N-butylacetamide (300

mg).

(+) APCI-MASS (m/z) : 637 (M+H)⁺

Example 66

5 under nitrogen, to a solution of N-benzyl-(3-hydroxy-
6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine
hydrochloride (300 mg) and (2S)-1-phenoxy-2,3-epoxypropane
(193 mg) in dichloromethane (10 ml) was added ytterbium(III)
trifluoromethanesulfonate (61 mg) at room temperature, and
10 the mixture was stirred at the same temperature for 3 hours.
The resulting mixture was poured into saturated aqueous
sodium bicarbonate solution and extracted with ethyl acetate.
The organic layer was washed with brine, dried over magnesium
sulfate and evaporated in vacuo. The residue was
15 chromatographed (hexane-ethyl acetate) over silica gel to
afford (2S)-1-[N-benzyl-N-(3-hydroxy-6,7,8,9-tetrahydro-5H-
benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol (340 mg).

NMR (CDCl₃, δ) : 1.20-1.40 (1H, m), 1.60-2.30 (4H, m),
2.50-2.95 (7H, m), 3.80-4.05 (5H, m), 6.50-6.75
20 (2H, m), 6.85-7.03 (4H, m), 7.20-7.38 (8H, m)

(+) APCI-MASS (m/z) : 418 (M+H)⁺

Example 67

25 Under nitrogen, to a solution of N-benzyl-(2,3-
dimethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine
(440 mg) and (2S)-1-phenoxy-2,3-epoxypropane (190 mg) in
dichloromethane (10 ml) was added ytterbium(III)
trifluoromethanesulfonate (157 mg) at room temperature, and
the mixture was stirred at the same temperature for 3 hours.
30 The resulting mixture was poured into saturated aqueous
sodium bicarbonate solution and extracted with ethyl acetate.
The organic layer was washed with brine, dried over magnesium
sulfate and evaporated in vacuo. The residue was
chromatographed (hexane-ethyl acetate) over silica gel to
35 afford (2S)-1-[N-benzyl-N-(2,3-dimethoxy-6,7,8,9-tetrahydro-

5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol (400 mg).

(+) APCI-MASS (m/z) : 462 (M+H)⁺

5 Example 68

Under nitrogen, to a solution of N-benzyl-(1,4-dimethoxy-5,6,8,9-tetrahydro-5H-benzocyclohepten-7-yl)amine (200 mg) and (2S)-1-phenoxy-2,3-epoxypropane (100 mg) in dichloromethane (10 ml) was added ytterbium(III) trifluoromethanesulfonate (80 mg) at room temperature, and the mixture was stirred at the same temperature for 3 hours. The resulting mixture was poured into saturated aqueous sodium bicarbonate solution and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford (2S)-1-[N-benzyl-N-(1,4-dimethoxy-5,6,8,9-tetrahydro-5H-benzocyclohepten-7-yl)amino]-3-phenoxy-2-propanol (70 mg).

(+) APCI-MASS (m/z) : 462 (M+H)⁺

20 Example 69

Under nitrogen, to a solution of N-benzyl-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-5-yl)amine (187 mg) and (2S)-1-phenoxy-2,3-epoxypropane (100 mg) in dichloromethane (10 ml) was added ytterbium(III) trifluoromethanesulfonate (50 mg) at room temperature, and the mixture was stirred at the same temperature for 3 hours. The resulting mixture was poured into saturated aqueous sodium bicarbonate solution and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford (2S)-1-[N-benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-5-yl)amino]-3-phenoxy-2-propanol (190 mg).

35 (+) APCI-MASS (m/z) : 432 (M+H)⁺

Example 70

To a solution of 1-[N-(3-aminomethyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)benzylamino]-3-phenoxy-2-propanol (150 mg) in N,N-dimethylformamide (2 ml) and 1N hydrochloric acid (0.8 ml) was added potassium cyanate (56.8 mg) at room temperature, and the mixture was stirred at the same temperature for 24 hours. The resulting mixture was poured into saturated aqueous sodium bicarbonate solution and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford N-[8-[N'-benzyl-N'-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylmethyl]urea (160 mg).

(+) APCI-MASS (m/z) : 474 (M+H)⁺

Example 71

To a solution of 1-[N-(3-aminomethyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)benzylamino]-3-phenoxy-2-propanol (140 mg) and triethylamine (0.046 ml) in tetrahydrofuran (10 ml) were added chloroformic acid 4-nitrophenyl ester (66.5 mg) and butylamine (0.030 ml) at room temperature, and the mixture was stirred at the same temperature for 18 hours. The resulting mixture was poured into saturated aqueous sodium bicarbonate solution and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford 1-[8-N-[benzyl-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylmethyl]-3-butylurea (110 mg).

(+) APCI-MASS (m/z) : 530 (M+H)⁺

Example 72

To a solution of (2S)-1-[N-(3-aminomethyl-6,7,8,9-

tetrahydro-5H-benzocyclohepten-6-yl)benzylamino]-3-phenoxy-2-propanol (100 mg) in methanol (2 ml) was added tert-butylacrylate (0.033 ml) at room temperature, and the mixture was stirred at the same temperature for 18 hours. The resulting mixture was evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford 3-[8-[N-benzyl-(2S)-(2-hydroxy-3-phenoxypropyl)amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylmethylamino]-propionic acid tert-butyl ester (100 mg).

(+) APCI-MASS (m/z) : 559 (M+H)⁺

Example 73

Under nitrogen, to a solution of 2-(8-benzylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy)-N-butylacetamide (170 mg) and (2S)-1-(4-benzyloxy-3-methanesulfonylamino)phenoxy-2,3-epoxypropane (108 mg) in dichloromethane (10 ml) was added ytterbium(III) trifluoromethanesulfonate (56 mg) at room temperature, and the mixture was stirred at the same temperature for 3 hours. The resulting mixture was poured into saturated aqueous sodium bicarbonate solution and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford 2-[8-[N-benzyl-N-[(2S)-3-(4-benzyloxy-3-methanesulfonylamino)phenoxy-2-hydroxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-butylacetamide (120 mg).

NMR (CDCl₃, δ) : 0.92 (3H, t, J=7.3Hz), 1.20-1.65 (7H, m), 1.95-2.05 (1H, m), 2.60-2.80 (4H, m), 2.60-2.80 (4H, m), 2.87 (3H, s), 3.30-3.40 (2H, m), 3.70-4.00 (5H, m), 4.45 (2H, d, J=2.0Hz), 5.05 (2H, s), 6.55-6.75 (2H, m), 6.80-7.25 (14H, m)

(+) APCI-MASS (m/z) : 730 (M+H)⁺

Example 74

Under nitrogen, to a solution of N-benzyl-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (487 mg) and (2S)-1-(4-benzyloxy-3-methanesulfonylamino)phenoxy-2,3-epoxypropane (338 mg) in dichloromethane (10 ml) was added ytterbium(III) trifluoromethanesulfonate (107 mg) at room temperature, and the mixture was stirred at the same temperature for 3 hours. The resulting mixture was poured into saturated aqueous sodium bicarbonate solution and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel to afford (2S)-1-[N-benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-(4-benzyloxy-3-methanesulfonylamino)phenoxy-2-propanol (750 mg).

(+) APCI-MASS (m/z) : 432 (M+H)⁺

Example 75

A mixture of (2S)-1-[N-benzyl-N-(6,7,8,9-tetrahydro-2-hydroxy-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol (300 mg) and 10% Pd/C (50% wet, 80 mg) in methanol (4.5 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 2 hours and 30 minutes and filtered. The filtrate was evaporated in vacuo, and the residue was chromatographed (chloroform-methanol) over silica gel (6.0 g). The eluate was treated with 4N hydrogen chloride in ethyl acetate and evaporated in vacuo. The residue was triturated in diethyl ether, and the precipitated powder was collected by decantation to afford (2S)-1-(6,7,8,9-tetrahydro-2-hydroxy-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol hydrochloride (238 mg) as a colorless powder.

mp : 53.5°C (dec.)

IR (KBr) : 3411, 2750-2650, 1241 cm⁻¹

NMR (DMSO- d_6 , δ) : 1.32 (1H, m), 1.75-2.05 (2H, m),
 2.24 (1H, m), 2.62 (2H, m), 2.95-3.45 (5H, m), 4.00
 (2H, m), 4.24 (1H, m), 5.90 (1H, d, $J=4.5\text{Hz}$), 6.50-
 6.55 (2H, m), 6.95-7.00 (4H, m), 7.31 (2H, t,
 5 $J=7.8\text{Hz}$), 8.77 (1H, br), 9.0 (1H, br), 9.28 (1H, s)

(+) APCI-MASS (m/z) : 328 (M+H)⁺

Anal. Calcd. for $\text{C}_{20}\text{H}_{26}\text{ClNO}_3 \cdot 0.5\text{H}_2\text{O}$:

C 64.42, H 7.30, N 3.76

Found : C 63.73, H 7.37, N 3.58

Example 76

The following compounds [Examples 76 and 77] were
 obtained according to a similar manner to that of Example 75.

(2S)-1-Phenoxy-3-(6,7,8,9-tetrahydro-1-hydroxy-5H-
 benzocyclohepten-6-yl)amino-2-propanol hydrochloride

mp : 73-84°C (dec.) (from diethyl ether)

IR (KBr) : 3357 (br), 2850-2650, 1242 cm^{-1}

NMR (DMSO- d_6 , δ) : 1.23 (2H, m), 1.75-2.05 (2H, m),
 20 2.28 (2H, m), 2.95-3.45 (5H, m), 4.00 (2H, m), 4.22
 (1H, m), 5.90 (1H, , $J=4.9\text{Hz}$), 6.63-6.75 (2H, m),
 6.87-7.00 (4H, m), 7.27-7.36 (2H, m), 8.74 (1H,
 br), 8.95 (1H, br), 9.34 (1H, s)

(+) APCI-MASS (m/z) : 328 (M+H)⁺

25 Anal. Calcd. for $\text{C}_{20}\text{H}_{26}\text{ClNO}_3 \cdot 1/4\text{H}_2\text{O}$:

C 65.21, H 7.25, N 3.80

Found : C 65.24, H 7.50, N 3.61

Example 77

30 1-(1,3-Dihydro-2-benzoimidazol-4-yl)oxy-3-(6,7,8,9-
 tetrahydro-2-hydroxy-5H-benzocyclohepten-6-yl)amino-2-
 propanol hydrochloride

mp : 100.5°C (dec.) (from diisopropyl ether)

IR (KBr) : 3410 (br), 3257 (br), 2800-2650, 1695, 1259,
 35 1244 cm^{-1}

NMR (DMSO- d_6 , δ) : 1.12-1.4 (1H, m), 1.75-2.2 (2H, m),
2.2-2.4 (1H, m), 2.4-2.8 (2H, m), 2.8-3.5 (5H, m),
3.95-4.3 (3H, m), 5.7 (1H, br), 6.51-6.66 (4H, m),
6.88 (1H, t, $J=8.2\text{Hz}$), 6.99 (1H, d, $J=6.9\text{Hz}$), 8.68
5 (1H, br), 8.93 (1H, br), 9.25 (1H, br)
(+) APCI-MASS (m/z) : 384 (M+H)⁺

Example 78

A mixture of (2S)-1-[N-benzyl-N-(3-chloro-6,7,8,9-
10 tetrahydro-2-hydroxy-5H-benzocyclohepten-6-yl)amino]-3-
phenoxy-2-propanol hydrochloride (121 mg) and 10% Pd/C (50%
wet, 18 mg) in chlorobenzene (1.8 ml) and MeOH (0.4 ml) was
stirred at room temperature in the presence of hydrogen at an
atmospheric pressure for 1 hour and filtered. The filtrate
15 was evaporated in vacuo, and the residue was partitioned
between ethyl acetate and a saturated aqueous solution of
sodium bicarbonate. The organic layer was separated, washed
with brine, dried over sodium sulfate and evaporated in
vacuo. The residue was chromatographed (dichloromethane-
20 methanol) over silica gel (2.0 g). The eluate was treated
with 4N hydrogen chloride in ethyl acetate and evaporated in
vacuo. The residue was triturated in diisopropyl ether, and
the precipitated powder was collected by filtration to afford
(2S)-1-(3-chloro-6,7,8,9-tetrahydro-2-hydroxy-5H-
25 benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol
hydrochloride (74 mg, 75%) as a colorless powder.

IR (KBr) : 3384 (br), 2800-2600, 1242 cm^{-1}

NMR (DMSO- d_6 , δ) : 1.17-1.31 (1H, m), 1.84-1.99 (2H,
m), 2.16-2.24 (1H, m), 2.61-2.74 (2H, m), 3.00-3.22
30 (4H, m), 3.98-4.24 (3H, m), 5.89 (1H, d, $J=3.9\text{Hz}$),
6.77 (1H, m), 6.86-7.02 (3H, m), 7.14-7.32 (3H, m),
8.73 (1H, br), 8.88 (1H, br), 10.03 (1H, s)

(+) APCI-MASS (m/z) : 362 and 364 (M+H)⁺

Example 79

A mixture of (2S)-1-[N-benzyl-N-(3-isopropoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol (281 mg) and 10% palladium on activated carbon (50% wet, 100 mg) in methanol (5 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 6 hours. After filtration, the filtrate was evaporated in vacuo, followed by treatment with 4N hydrogen chloride in ethyl acetate to give (2S)-1-(3-isopropoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol hydrochloride (130 mg).

(+) APCI-MASS (m/z) : 370 (M-HCl+H)⁺

The following compounds [Examples 80 to 108] were obtained according to a similar manner to that of Example 79.

Example 80

(2S)-1-(3-Hexyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol hydrochloride

NMR (DMSO-d₆, δ) : 0.8 (3H, m), 1.1-1.5 (7H, m), 1.6-2.15 (4H, m), 2.25-2.4 (1H, m), 2.5-2.8 (2H, m), 2.9-3.5 (5H, m), 3.85-4.1 (4H, m), 4.15-4.25 (1H, m), 5.9-6.05 (1H, m), 6.65-7.1 (6H, m), 7.2-7.4 (2H, m), 8.7-9.3 (2H, m)

(+) APCI-MASS (m/z) : 412 (M+H)⁺

Example 81

(2S)-1-[3-(3-Cyano-pyridin-2-yl)oxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]amino]-3-phenoxy-2-propanol hydrochloride

(+) ESI-MASS (m/z) : 464 (M-HCl-H)⁺

Example 82

(2S)-1-[3-(2-Methoxyethoxy)-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]amino-3-phenoxy-2-propanol hydrochloride

(+) APCI-MASS (m/z) : 386 (M-HCl+H)⁺

Example 83

5 N-(2-Dimethylaminoethyl)-2-[8-[(2S)-2-hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetamide dihydrochloride

(+) APCI-MASS (m/z) : 456 (M-HCl+H)⁺

10 Example 84

(2S)-1-(3-Amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol dihydrochloride

(+) APCI-MASS (m/z) : 327 (M-HCl+H)⁺

15 Example 85

(2S)-1-(3-Dimethylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol hydrochloride

(+) APCI-MASS (m/z) : 355 (M-HCl+H)⁺

20 Example 86

(2S)-1-(3-Dimethylamino-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol hydrochloride

(+) APCI-MASS (m/z) : 355 (M-HCl+H)⁺

25 Example 87

N-[8-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yl]acetamide hydrochloride

(+) ESI-MASS (m/z) : 369 (M-HCl+H)⁺

30 Example 88

N-[8-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yl]benzamide hydrochloride

(+) APCI-MASS (m/z) : 431 (M-HCl+H)⁺

35 Example 89

N-[8-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yl]methanesulfonamide hydrochloride

(+) APCI-MASS (m/z) : 405 (M-HCl+H)⁺

5

Example 90

N-[8-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yl]benzenesulfonamide hydrochloride

10 (+) APCI-MASS (m/z) : 467 (M-HCl+H)⁺

Example 91

6-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-3-methoxycarbonylamino-6,7,8,9-tetrahydro-5H-benzocycloheptene hydrochloride

15

(+) APCI-MASS (m/z) : 385 (M-HCl+H)⁺

Example 92

6-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-3-phenoxy carbonylamino-6,7,8,9-tetrahydro-5H-benzocycloheptene hydrochloride

20

(+) APCI-MASS (m/z) : 447 (M-HCl+H)⁺

Example 93

N,N-Dimethyl-N'-[8-[(2S)-2-hydroxy-3-phenoxypropyl]-amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yl]sulfonamide hydrochloride

25

(+) APCI-MASS (m/z) : 434 (M-HCl+H)⁺

30 Example 94

N-[8-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yl]sulfonamide hydrochloride

(+) APCI-MASS (m/z) : 406 (M-HCl+H)⁺

35 Example 95

(2S)-3-Phenoxy-1-(6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-2-propanol hydrochloride

(+) ESI-MASS (m/z) : 312 (M-HCl+H)⁺

5 Example 96

(2S)-1-Phenoxy-3-(3-phenyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-2-propanol hydrochloride

(+) ESI-MASS (m/z) : 388 (M-HCl+H)⁺

10 Example 97

(2S)-3-Phenylsulfonyl-1-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-2-propanol hydrochloride

(+) APCI-MASS (m/z) : 390 (M-HCl+H)⁺

15 Example 98

1-(3-Methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenylamino-2-propanol dihydrochloride

(+) APCI-MASS (m/z) : 341 (M-2HCl+H)⁺

20 Example 99

(2S)-1-(6-Aminopyridin-3-yl)oxy-3-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-2-propanol trihydrochloride

(+) APCI-MASS (m/z) : 358 (M-3HCl+H)⁺

25

Example 100

(2S)-1-(4-Fluorophenoxy)-3-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-2-propanol hydrochloride

30

NMR (DMSO-d₆, δ) : 1.1-1.4 (1H, m), 1.7-2.1 (2H, m),
2.2-2.4 (1H, m), 2.6-2.8 (2H, m), 2.95-3.5 (5H, m),
3.72 (3H, s), 3.9-4.0 (2H, m), 4.22 (1H, br s),
5.93 (1H, br s), 6.71 (1H, dd, J=2.6 and 8.2Hz),
6.82 (1H, dd, J=2.6 and 6.5Hz), 6.9-7.25 (5H, m),
8.82 (1H, br s), 9.16 (1H, br s)

35

(+) APCI-MASS (m/z) : 360 (M-HCl+H)⁺

Example 101

(2S)-1-(4-Methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol hydrochloride

IR (KBr) : 3369, 2937, 1591, 1452, 1259, 1236,
1074 cm⁻¹

NMR (DMSO-d₆, δ) : 1.22-1.42 (1H, m), 1.80-2.10 (2H, m), 2.25-2.39 (1H, m), 2.66-2.83 (3H, m), 2.90-3.23 (3H, m), 3.53-3.60 (1H, m), 3.78 (3H, s), 3.98-4.05 (2H, m), 4.20-4.31 (1H, m), 5.91 (1H, br), 6.75 (1H, d, J=7.9Hz), 6.86 (1H, d, J=7.9Hz), 6.92-7.00 (3H, m), 7.12 (1H, dd, J=7.9 and 7.9Hz), 7.32 (2H, dd, J=8.1 and 8.1Hz), 8.89 (2H, br)

(+) APCI-MASS (m/z) : 342 (M+H)⁺

Example 102

8-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxylic acid methyl ester hydrochloride

IR (KBr) : 3378, 2950, 2796, 1716, 1438, 1280, 1243 cm⁻¹

NMR (DMSO-d₆, δ) : 1.20-1.42 (1H, m), 1.81-2.13 (2H, m), 2.25-2.40 (1H, m), 2.74-2.91 (2H, m), 3.02-3.37 (5H, m), 3.84 (3H, s), 4.01 (2H, d, J=4.6Hz), 4.20-4.36 (1H, m), 5.93 (1H, br), 6.92-6.99 (3H, m), 7.27-7.35 (3H, m), 7.76 (1H, d, J=7.8Hz), 7.87 (1H, d, J=7.8Hz), 8.92-9.18 (2H, br)

(+) ESI-MASS (m/z) : 370 (M+H)⁺

Example 103

8-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxamide hydrochloride

IR (KBr) : 3353, 2937, 1662, 1600, 1243 cm⁻¹

NMR (DMSO-d₆, δ) : 1.21-1.42 (1H, m), 1.78-2.13 (2H, m), 2.24-2.40 (1H, m), 2.74-2.85 (2H, m), 3.02-3.44

(5H, m), 4.01 (2H, d, J=4.5Hz), 4.18-4.30 (1H, m),
 5.92 (1H, d, J=4.0Hz), 6.92-6.99 (3H, m), 7.21 (1H,
 d, J=7.8Hz), 7.27-7.35 (3H, m), 7.67 (1H, d,
 J=7.8Hz), 7.79 (1H, d, J=2.0Hz), 7.91 (1H, s), 8.84
 (1H, br s), 9.12 (1H, br s)

(+) APCI-MASS (m/z) : 355 (M+H)⁺

Example 104

N-[8-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-6,7,8,9-
 tetrahydro-5H-benzocyclohepten-2-ylmethyl]acetamide
 hydrochloride

IR (KBr) : 3278, 2935, 1648, 1494, 1241 cm⁻¹

NMR (DMSO-d₆, δ) : 1.17-1.37 (1H, m), 1.78-2.10 (2H,
 m), 1.86 (3H, s), 2.25-2.37 (1H, m), 2.67-2.77 (2H,
 m), 2.98-3.40 (5H, m), 4.01 (2H, d, J=3.9Hz), 4.15-
 4.35 (1H, m), 4.19 (2H, d, J=5.7Hz), 6.92-7.10 (6H,
 m), 7.31 (2H, dd, J=7.8 and 7.8Hz), 8.30-8.40 (1H,
 m), 8.85 (1H, br s), 9.20 (1H, br s)

(+) APCI-MASS (m/z) : 383 (M+H)⁺

Example 105

N-[8-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-6,7,8,9-
 tetrahydro-5H-benzocyclohepten-2-ylmethyl]benzamide
 hydrochloride

IR (KBr) : 3330, 2933, 1639, 1538, 1455, 1241 cm⁻¹

NMR (DMSO-d₆, δ) : 1.18-1.38 (1H, m), 1.79-2.08 (2H,
 m), 2.23-2.37 (1H, m), 2.68-2.78 (2H, m), 2.98-3.30
 (5H, m), 3.97-4.03 (2H, m), 4.16-4.28 (1H, m), 4.44
 (2H, d, J=5.4Hz), 5.91 (1H, br), 6.91-6.98 (3H, m),
 7.10-7.18 (3H, m), 7.30 (2H, dd, J=7.7 and 7.7Hz),
 7.41-7.57 (3H, m), 7.91 (2H, d, J=7.7Hz), 8.75-9.10
 (2H, br), 9.03-9.13 (1H, m)

(+) APCI-MASS (m/z) : 445 (M+H)⁺

Example 106

N-[8-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylmethyl]methanesulfonamide hydrochloride

IR (KBr) : 3326, 2933, 1594, 1496, 1313, 1241, 1147 cm^{-1}

5 NMR (DMSO- d_6 , δ) : 1.15-1.38 (1H, m), 1.75-2.09 (2H, m), 2.23-2.37 (1H, m), 2.68-2.78 (2H, m), 2.87 (3H, s), 2.93-3.32 (5H, m), 4.00 (2H, d, $J=4.1\text{Hz}$), 4.10 (2H, d, $J=6.3\text{Hz}$), 4.17-4.30 (1H, m), 5.93 (1H, br), 6.91-7.00 (3H, m), 7.11-7.20 (3H, m), 7.31 (2H, dd, $J=7.8$ and 7.8Hz), 7.50-7.59 (1H, m), 8.80-9.10 (2H, br)

(+) APCI-MASS (m/z) : 419 (M+H)⁺

Example 107

15 N-[8-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylmethyl]benzenesulfonamide hydrochloride

IR (KBr) : 2935, 1594, 1496, 1448, 1322, 1241, 1157 cm^{-1}

20 NMR (DMSO- d_6 , δ) : 1.13-1.38 (1H, m), 1.75-2.08 (2H, m), 2.24-2.38 (1H, m), 2.65-2.75 (2H, m), 2.92-3.35 (5H, m), 3.92 (2H, d, $J=6.2\text{Hz}$), 4.01 (2H, d, $J=4.4\text{Hz}$), 4.18-4.30 (1H, m), 5.92 (1H, br), 6.92-7.02 (5H, m), 7.10 (1H, s), 7.31 (2H, dd, $J=7.8$ and 7.8Hz), 7.50-7.68 (3H, m), 7.79 (2H, d, $J=7.8\text{Hz}$), 8.12-8.22 (1H, m), 8.86 (2H, br)

25 (+) APCI-MASS (m/z) : 481 (M+H)⁺

Example 108

30 1-(3-Methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-4-phenyl-2-butanol hydrochloride

(+) APCI-MASS (m/z) : 340 (M-HCl+H)⁺

Example 109

35 A solution of N-(3-ethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-

(triethylsilyloxy)propyl]-tert-butoxycarbonylamine (115 mg) and concentrated hydrochloric acid (0.5 ml) in methanol (5 ml) was stirred at room temperature for 48 hours. The mixture was evaporated in vacuo. The residue was triturated with diethyl ether-diisopropyl ether, followed by decantation and dryness in vacuo to give (2S)-1-(3-ethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol hydrochloride (30 mg).

NMR (DMSO- d_6 , δ) : 1.1-1.4 (4H, m), 1.7-2.1 (2H, m), 2.2-2.35 (1H, m), 2.55-2.75 (2H, m), 2.9-3.4 (5H, m), 3.9-4.1 (4H, m), 4.25 (1H, br s), 5.9-6.0 (1H, m), 6.69 (1H, dd, $J=2.5$ and 8.2Hz), 6.82 (1H, dd, $J=2.4$ and 6.8Hz), 6.9-7.1 (4H, m), 7.31 (2H, t, $J=7.9\text{Hz}$), 8.7-9.4 (2H, m)

(+) APCI-MASS (m/z) : 356 (M+H)⁺

Example 110

To N-(3-isopropoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-butoxycarbonylamine (79 mg) was added 4N hydrogen chloride in ethyl acetate (3 ml) at room temperature, and the mixture was stirred at room temperature for 1 hour. After evaporation in vacuo, the residue was triturated with diisopropyl ether and dried in vacuo to give (2S)-1-(3-isopropoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol hydrochloride (35 mg).

(+) APCI-MASS (m/z) : 370 (M-HCl+H)⁺

The following compounds [Examples 111 to 116] were obtained according to a similar manner to that of Example 110.

Example 111

(2S)-1-(3-Allyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol hydrochloride

(-) ESI-MASS (m/z) : 402 (M-HCl-H)⁺

Example 112

(2S)-1-(3-Cyclopentyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol hydrochloride

(+) APCI-MASS (m/z) : 396 (M-HCl+H)⁺

Example 113

(2S)-1-(3-Benzylloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol hydrochloride

(+) APCI-MASS (m/z) : 418 (M-HCl+H)⁺

Example 114

(2S)-1-(3-Phenethyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol hydrochloride

(+) APCI-MASS (m/z) : 432 (M-HCl+H)⁺

Example 115

(2S)-1-(3-Carbamoyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol hydrochloride

(+) APCI-MASS (m/z) : 371 (M-HCl+H)⁺

Example 116

(2S)-1-(3-Dimethylsulfonyloxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol

NMR (DMSO-d₆, δ) : 1.2-1.45 (1H, m), 1.7-2.1 (2H, m),
2.3-2.4 (1H, m), 2.7-3.35 (7H, m), 2.90 (6H, m),
4.00 (2H, d, J=8.6Hz), 4.2-4.35 (1H, m), 6.9-7.35
(8H, m)

(+) APCI-MASS (m/z) : 435 (M-HCl+H)⁺

Example 117

To a solution of N-(3-fluoromethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-2-hydroxy-3-phenoxypropyl]-

tert-butoxycarbonylamine (41 mg) in ethyl acetate (2 ml) was added 4N hydrogen chloride in ethyl acetate (0.45 ml) at room temperature, and the mixture was stirred at the same temperature for 5.5 hours. The mixture was poured into a mixture of saturated aqueous sodium hydrogencarbonate and ethyl acetate, followed by being made basic with saturated aqueous sodium hydrogencarbonate. After separation, the organic layer was dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by preparative thin layer chromatography (silica gel, chloroform:methanol = 10:1) to give (2S)-1-(3-fluoromethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol (23 mg).

(+) APCI-MASS (m/z) : 360 (M+H)⁺

Example 118

A mixture of 2-[8-[N-benzyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetamide methanesulfonate (240 mg) and 10% palladium on activated carbon (50% wet, 50 mg) in ethanol (5 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 2 hours. After filtration, the filtrate was evaporated in vacuo and followed by trituration with diisopropyl ether to give 2-[8-[(2S)-2-hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetamide methanesulfonate (187 mg).

(+) APCI-MASS (m/z) : 385 (M-MsOH+H)⁺

Example 119

A solution of 2-[8-[N-benzyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid ethyl ester (200 mg) and ethylamine (70% in water, 77 μ l) in methanol (2 ml) were stirred at room temperature for 12 hours. After evaporation in vacuo, the crude product and 10% palladium on activated carbon (50% wet, 60 mg) in methanol (2 ml) were stirred at room temperature in

the presence of hydrogen at an atmospheric pressure for 2.5 hours. After filtration, the filtrate was evaporated in vacuo. The residue was purified by column chromatography on silica gel (chloroform:methanol = 10:1) and followed by treatment with 4N hydrogen chloride in ethyl acetate to give N-ethyl-2-[8-[(2S)-2-hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetamide hydrochloride (120 mg).

NMR (DMSO- d_6 , δ) : 1.0-1.1 (3H, m), 1.1-1.4 (1H, m), 1.7-2.1 (2H, m), 2.25-2.4 (1H, m), 2.6-2.8 (2H, m), 2.9-3.7 (7H, m), 3.95-4.1 (2H, m), 4.15-4.35 (1H, m), 4.40 (2H, s), 6.73 (1H, dd, J=2.6 and 8.3Hz), 6.85-7.1 (5H, m), 7.31 (2H, t, J=8.0Hz), 8.15-8.2 (1H, m), 8.7-9.3 (2H, m)

Example 120

A solution of 2-[8-[N-benzyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid ethyl ester (220 mg) and n-hexylamine (173 μ l) in methanol (2 ml) was stirred at room temperature for 36 hours. And then to this one was added 10% palladium on activated carbon (50% wet, 60 mg), and the mixture was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 4 hours. After filtration, the filtrate was evaporated in vacuo. The residue was purified by preparative thin layer chromatography (silica gel, chloroform:methanol = 10:1) and followed by treatment with 4N hydrogen chloride in ethyl acetate to give N-hexyl-2-[8-[(2S)-2-hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetamide hydrochloride (169 mg).

(+) APCI-MASS (m/z) : 469 (M-HCl+H)⁺

The following compound was obtained according to a similar manner to that of Example 120.

Example 121

2-[8-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-(2-methoxyethyl)-acetamide hydrochloride

5 NMR (DMSO-d₆, δ) : 1.1-1.4 (1H, m), 1.7-2.1 (2H, m),
2.25-2.4 (1H, m), 2.6-2.8 (2H, m), 3.0-3.45 (12H,
m), 3.95-4.1 (2H, m), 4.2-4.35 (1H, m), 4.43 (2H,
s), 5.9-6.0 (1H, m), 6.7-6.8 (1H, m), 6.85-7.1 (5H,
m), 7.25-7.4 (2H, m), 8.0-8.1 (1H, m), 8.7-9.3 (2H,
10 m)

(+) APCI-MASS (m/z) : 443 (M-HCl+H)⁺

Example 122

15 To a solution of N-[3-(N'-butylmethylcarbamoyl)methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]-N-[(2S)-2-hydroxy-3-phenoxypropyl]-tert-butoxycarbonylamine (85 mg) in ethyl acetate (2 ml) was added 4N hydrogen chloride in ethyl acetate (1 ml) at 5°C, and the mixture was stirred at room temperature for 2 hours. After evaporation in vacuo, the residue was triturated with diisopropyl ether and dried in vacuo to give N-butyl-2-[8-[(2S)-2-hydroxy-3-phenoxypropyl]-amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-methylacetamide hydrochloride (45 mg).

20 NMR (DMSO-d₆, δ) : 0.88 (3H, q, J=7.3Hz), 1.10-1.65
25 (5H, m), 1.70-2.16 (2H, m), 2.23-2.40 (1H, m),
2.58-3.43 (12H, m), 3.85-4.10 (2H, m), 4.17-4.35
(1H, m), 4.73 (2H, s), 5.83-5.99 (1H, m), 6.59-7.08
(6H, m), 7.31 (2H, t, J=7.7Hz), 8.81 (1H, br s),
9.15 (1H, br s)

30 (+) APCI-MASS (m/z) : 455 (M+H)⁺

The following compounds [Examples 123 to 128] were obtained according to a similar manner to that of Example 122.

Example 123

N-sec-Butyl-2-[8-[(2S)-2-hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetamide hydrochloride

5 (+) APCI-MASS (m/z) : 441 (M+H)⁺

Example 124

N-Cyclohexyl-2-[8-[(2S)-2-hydroxy-3-phenoxypropyl]-amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetamide hydrochloride

10 (+) APCI-MASS (m/z) : 467 (M+H)⁺

Example 125

N-Benzyl-2-[8-[(2S)-2-hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetamide hydrochloride

15 (+) APCI-MASS (m/z) : 475 (M+H)⁺

Example 126

20 2-[8-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-(2-methylthioethyl)acetamide hydrochloride

 (+) APCI-MASS (m/z) : 459 (M+H)⁺

Example 127

25 6-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-3-piperidinocarbonylmethoxy-6,7,8,9-tetrahydro-5H-benzocycloheptene hydrochloride

 (+) APCI-MASS (m/z) : 453 (M+H)⁺

Example 128

30 2-[8-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-(1H-indol-5-yl)acetamide hydrochloride

35 NMR (DMSO-d₆-δ) : 1.14-1.49 (1H, m), 1.70-2.12 (2H, m),

2.25-2.39 (1H, m), 2.57-2.82 (2H, m), 2.94-4.13
(7H, m), 4.19-4.32 (1H, m), 4.52-4.78 (2H, m),
6.72-7.49 (13H, m), 7.50-7.93 (1H, m), 8.69-9.33

5 (+) APCI-MASS (m/z) : 500 (M+H)⁺

Example 129

Under nitrogen, to a solution of (2S)-1-amino-3-phenoxy-
2-propanol (280 mg) in 1,2-dichloroethane (10 ml) were added
10 3-chloro-5,7,8,9-tetrahydrobenzocyclohepten-6-one (300 mg),
sodium triacetoxymethylborohydride (820 mg) and acetic acid (0.26
ml) at room temperature, and the mixture was stirred at the
same temperature for 6.5 hours. The resulting mixture was
poured into aqueous 1N sodium hydroxide and extracted with
15 ethyl acetate. The organic layer was washed with brine,
dried over anhydrous magnesium sulfate and evaporated in
vacuo. The residue was purified by column chromatography on
silica gel (chloroform:methanol = 20:1 to 10:1) and followed
by treatment with 4N hydrogen chloride in ethyl acetate to
20 give (2S)-1-(3-chloro-6,7,8,9-tetrahydro-5H-benzocyclohepten-
6-
yl)amino-3-phenoxy-2-propanol hydrochloride (260 mg).

(+) ESI-MASS (m/z) : 346, 348 (M-HCl+H)⁺

25 The following compounds [Examples 130 and 131] were
obtained according to a similar manner to that of Example
129.

Example 130

30 (2S)-1-(3-Nitro-6,7,8,9-tetrahydro-5H-benzocyclohepten-
6-yl)amino-3-phenoxy-2-propanol hydrochloride
(+) ESI-MASS (m/z) : 357 (M-HCl+H)⁺

Example 131

35 (2S)-1-(3-Methyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-

6-yl)amino-3-phenoxy-2-propanol hydrochloride

(+) ESI-MASS (m/z) : 325 (M-HCl+H)⁺

Example 132

5 A mixture of (2S)-1-[N-benzyl-N-(2-chloro-3-methoxy-
6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-
2-propanol (172 mg) and 10% palladium on activated carbon
(50% wet, 50 mg) in a mixture of chlorobenzene (2 ml) and
methanol (2 ml) was stirred at room temperature in the
10 presence of hydrogen at an atmospheric pressure for 2 hours.
After filtration, the filtrate was evaporated in vacuo. The
residue was purified by column chromatography on silica gel
(chloroform:methanol = 20:1 to 10:1) and followed by
treatment with 4N hydrogen chloride in ethyl acetate to give
15 1-(2-chloro-3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-
6-yl)amino-3-phenoxy-2-propanol hydrochloride (95 mg).

(+) APCI-MASS (m/z) : 376, 378 (M-HCl+H)⁺

Example 133

20 Under nitrogen, to a solution of 3-methoxy-6,7,8,9-
tetrahydro-5H-benzocyclohepten-6-ylamine (330 mg) and (2S)-1-
phenylthio-2,3-epoxypropane (290 mg) in dichloromethane (5
ml) was added ytterbium(III) trifluoromethanesulfonate (110
mg) at room temperature, and the mixture was stirred at the
25 same temperature for 24 hours. The resulting mixture was
poured into saturated aqueous sodium hydrogencarbonate and
extracted with ethyl acetate. The organic layer was washed
with brine, dried over anhydrous magnesium sulfate and
evaporated in vacuo. The residue was purified by column
30 chromatography on silica gel (chloroform:methanol = 50:1 to
20:1) followed by treatment with 4N hydrogen chloride in
ethyl acetate to give (2S)-1-(3-methoxy-6,7,8,9-tetrahydro-
5H-benzocyclohepten-6-yl)amino-3-phenylthio-2-propanol
hydrochloride (360 mg).

35 (+) APCI-MASS (m/z) : 358 (M-HCl+H)⁺

Example 134

8-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxylic acid methyl ester hydrochloride was converted to the corresponding free base in a usual manner. A solution of the base (91 mg) in methanol (1 ml) and 1N sodium hydroxide aqueous solution (0.246 ml) was stirred at 70°C for 6 hours and evaporated in vacuo. The residue was triturated in diisopropyl ether, and the precipitated powder was collected by filtration to afford sodium 8-[(2S)-2-hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocycloheptene-2-carboxylate (74 mg).

IR (KBr) : 3380, 2925, 1592, 1550, 1386, 1245 cm^{-1}

NMR (DMSO-d_6 , δ) : 1.23-1.70 (3H, m), 1.75-2.06 (2H, m), 2.38-2.90 (7H, m), 3.79-3.98 (3H, m), 5.11 (1H, br), 6.87-6.98 (4H, m), 7.27 (2H, dd, $J=8.0$ and 8.0Hz), 7.58 (1H, d, $J=7.5\text{Hz}$), 7.65 (1H, s)

(+) ESI-MASS (m/z) : 356 ($\text{M-Na}+2\text{H}$)⁺, 378 ($\text{M}+\text{H}$)⁺

Example 135

A solution of N-[(2S)-2-hydroxy-3-phenoxypropyl]-N-(3-methylcarbamoyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-tert-butoxycarbonylamine (108 mg) in ethyl acetate (1 ml) was treated with 4N hydrogen chloride in ethyl acetate solution (0.58 ml) overnight. The mixture was evaporated in vacuo and triturated in diisopropyl ether, and the precipitated powder was collected by filtration to afford 6-[(2S)-2-hydroxy-3-phenoxypropyl]amino-3-methylcarbamoyl-6,7,8,9-tetrahydro-5H-benzocycloheptene hydrochloride (86 mg).

IR (KBr) : 3332, 2937, 1631, 1457, 1241 cm^{-1}

NMR (DMSO-d_6 , δ) : 1.22-1.42 (1H, m), 1.80-2.11 (2H, m), 2.23-2.39 (1H, m), 2.77 (3H, d, $J=4.4\text{Hz}$), 2.79-2.86 (2H, m), 3.02-3.45 (5H, m), 4.00 (2H, d, $J=4.0\text{Hz}$), 4.19-4.30 (1H, m), 5.83 (1H, br), 6.92-6.99 (3H, m), 7.21 (1H, d, $J=7.8\text{Hz}$), 7.31 (2H, dd, $J=8.0$ and 8.0Hz), 7.62 (1H, d, $J=7.8\text{Hz}$), 7.75 (1H,

s), 8.40 (1H, t like, $J=4.4\text{Hz}$), 8.81 (1H, br s),
9.09 (1H, br s)

(+) ESI-MASS (m/z) : 369 ($M+H$)⁺

5 The following compounds [Examples 136 and 137] were
obtained according to a similar mannder to that of Example
135.

Example 136

10 6-[(2S)-2-Hydroxy-3-phenoxypropyl]amino-3-
dimethylcarbamoyl-6,7,8,9-tetrahydro-5H-benzocycloheptene
hydrochloride

IR (KBr) : 3382, 2935, 1602, 1492, 1241 cm^{-1}

15 NMR (DMSO- d_6 , δ) : 1.22-1.43 (1H, m), 1.83-2.10 (2H,
m), 2.23-2.38 (1H, m), 2.76-2.83 (2H, m), 2.94 (6H,
s), 3.00-3.40 (5H, m), 3.97-4.03 (2H, m), 4.18-4.30
(1H, m), 6.90-7.02 (3H, m), 7.19-7.35 (5H, m), 8.80
(1H, br), 9.10 (1H, br)

(+) ESI-MASS (m/z) : 383 ($M+H$)⁺

Example 137

25 6-[(2S)-2-Hydroxy-3-phenylpropyl]amino-3-
phenylcarbamoyl-6,7,8,9-tetrahydro-5H-benzocycloheptene
hydrochloride

IR (KBr) : 3380, 2937, 1648, 1598, 1536, 1498, 1241 cm^{-1}

30 NMR (DMSO- d_6 , δ) : 1.20-1.44 (1H, m), 1.80-2.18 (2H,
m), 2.20-2.40 (1H, m), 2.78-2.90 (2H, m), 3.02-3.40
(5H, m), 3.98-4.08 (2H, m), 4.18-4.31 (1H, m), 5.92
(1H, d, $J=4.9\text{Hz}$), 6.91-6.99 (4H, m), 7.09 (1H, dd,
 $J=7.3$ and 7.3Hz), 7.26-7.39 (5H, m), 7.75-7.87 (4H,
m), 8.76 (1H, br), 8.91 (1H, br), 10.21 (1H, s)

(+) ESI-MASS (m/z) : 431 ($M+H$)⁺

Example 138

35 To N-(3-hydroxymethyl-6,7,8,9-tetrahydro-5H-

benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-butoxycarbonylamine (200 mg) was added 4N hydrogen chloride in ethyl acetate (6 ml) at room temperature, and the mixture was stirred at the same temperature for 1 hour. After evaporation in vacuo, the residue was dissolved into a mixture of saturated aqueous sodium hydrogencarbonate and ethyl acetate. After separation, the organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by preparative thin layer chromatography (silica gel, chloroform:methanol = 5:1) and followed by treatment with 4N hydrogen chloride in ethyl acetate to give (2S)-1-(3-hydroxymethyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol hydrochloride (45 mg).

(+) APCI-MASS (m/z) : 342 (M-HCl+H)⁺

Example 139

A solution of N-(3-hydroxymethyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-N-[(2S)-3-phenoxy-2-(triethylsilyloxy)propyl]-tert-butoxycarbonylamine (188 mg) in ethyl acetate (2 ml) was treated with 4N hydrogen chloride in ethyl acetate (2 ml) for 3 hours. The mixture was evaporated in vacuo and washed with diethyl ether three times. The residue was triturated in diisopropyl ether, and precipitated powder was collected by filtration to afford 6-[(2S)-2-hydroxy-3-phenoxypropyl]amino-3-acetyloxymethyl-6,7,8,9-tetrahydro-5H-benzocycloheptene (113 mg).

NMR (DMSO-d₆, δ) : 1.12-1.41 (1H, m), 1.75-2.10 (2H, m), 2.05 (3H, s), 2.22-2.39 (1H, m), 2.65-2.85 (2H, m), 2.97-3.37 (5H, m), 3.94-4.04 (2H, m), 4.18-4.35 (1H, m), 5.02 (2H, s), 5.93 (1H, d, J=4.3Hz), 6.92-7.00 (3H, m), 7.08-7.38 (5H, m), 8.89 (1H, br), 9.27 (1H, br)

(+) APCI-MASS (m/z) : 384 (M+H)⁺

The following compound was obtained according to a similar manner to that of Example 139.

Example 140

(2S)-1-(3-Dimethylaminomethyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-3-phenoxy-2-propanol dichloride

IR (KBr) : 3380, 2937, 2709, 1594, 1494, 1457,
1243 cm^{-1}

NMR (DMSO- d_6 , δ) : 1.12-1.43 (1H, m), 1.80-2.10 (2H, m), 2.22-2.42 (1H, m), 2.67 (6H, s), 2.72-2.88 (2H, m), 2.98-3.45 (5H, m), 3.95-4.05 (2H, m), 4.10-4.35 (3H, m), 5.92 (1H, br), 6.90-7.02 (3H, m), 7.16-7.42 (5H, m), 8.87 (1H, br), 9.27 (1H, br), 10.81 (1H, br)

(+) APCI-MASS (m/z) : 369 (M+H)⁺

Example 141

A mixture of N-[8-[(2S)-2-hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yl]methyl acetamide hydrochloride (100 mg) and 1N aqueous hydrochloric acid solution (2 ml) was heated to reflux for 18 hours. The solvent was removed under reduced pressure to dryness. The residue was triturated in diisopropyl ether, and the precipitated powder was collected by filtration to afford (2S)-1-[(3-aminomethyl-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol dihydrochloride (94 mg).

IR (KBr) : 3382, 2933, 1594, 1494, 1241 cm^{-1}

NMR (DMSO- d_6 , δ) : 1.12-1.38 (1H, m), 1.77-2.10 (2H, m), 2.28-2.42 (1H, m), 2.70-2.82 (2H, m), 2.95-3.32 (5H, m), 3.88-4.00 (2H, m), 4.02 (2H, d, J=4.6Hz), 4.23-4.33 (1H, m), 5.94 (1H, br), 6.92-6.99 (3H, m), 7.18 (1H, d, J=7.5Hz), 7.27-7.74 (4H, m), 8.49 (3H, br s), 9.00 (1H, br s), 9.46 (1H, br s)

(+) APCI-MASS (m/z) : 341 (M+H)⁺

Example 142

A mixture of N-[(2R)-3-(3-methylsulfonylamino-4-benzyloxy)phenyl-2-hydroxypropyl]-N-[(6RS)-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)benzylamine (34 mg), 50% wet palladium (10%) on charcoal (10 mg) and methanol (1 ml) was stirred under hydrogen (1 atm) at room temperature for 1.5 hours. The reaction mixture was filtered and evaporated. The residue was purified by preparative TLC (eluent; 25% methanol-dichloromethane) to give N-[(2R)-3-(3-methylsulfonylamino-4-hydroxy)phenyl-2-hydroxypropyl]-[(6RS)-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (6.9 mg).

IR (KBr) : 3446, 2925, 1612, 1500, 1267, 1157, 1106 cm^{-1}

NMR (CD_3OD , δ) : 1.4-2.2 (4H, m), 2.40-2.9 (9H, m), 2.86 (3H, d, $J=2.0\text{Hz}$), 3.74 (3H, s), 3.7-3.9 (1H, m), 6.57-6.83 (4H, m), 6.95 (1H, d, $J=8.1\text{Hz}$), 7.15 (1H, br s)

MASS (m/z) : 435 (M)⁺

Example 143

A mixture of (2S)-3-[N-benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-1-(4-acetylamino)phenoxy-2-propanol (80 mg) and 10% palladium on activated carbon (50% wet, 10 mg) in methanol (10 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 3 hours and filtered. The filtrate was evaporated in vacuo and treated with 4N hydrogen chloride in ethyl acetate to afford (2S)-3-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-1-(4-acetylamino)-phenoxy-2-propanol hydrochloride (80 mg).

(+) APCI-MASS (m/z) : 435 ($\text{M}+\text{H}$)⁺

Example 144

A mixture of 2-[8-[N'-benzyl-[(2S)-2-hydroxy-3-(1H-

indol-5-yloxy)propyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yl]oxy-N-butylacetamide (160 mg) and 10% palladium on activated carbon (50% wet, 30 mg) in methanol (10 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 3 hours and filtered. The filtrate was evaporated in vacuo and treated with 4N hydrogen chloride in ethyl acetate to afford N-butyl-2-[8-[(2S)-2-hydroxy-3-(1H-indol-5-yloxy)propylamino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetamide hydrochloride (133 mg).

(+) APCI-MASS (m/z) : 480 (M+H)⁺

Example 145

A mixture of 2-[8-[N'-benzyl-N'-[(2S)-3-(4-benzyloxyphenoxy)-2-hydroxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yl]oxy-N-butylacetamide (300 mg) and 10% palladium on activated carbon (50% wet, 100 mg) in methanol (10 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 3 hours and filtered. The filtrate was evaporated in vacuo and treated with 4N hydrogen chloride in ethyl acetate to afford N-butyl-2-[8-[(2S)-2-hydroxy-3-(4-hydroxyphenoxy)propyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetamide hydrochloride (190 mg).

(+) APCI-MASS (m/z) : 457 (M+H)⁺

Example 146

A mixture of (2S)-1-[N-benzyl-N-(3-hydroxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol (320 mg) and 10% palladium on activated carbon (50% wet, 100 mg) in methanol (20 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 3 hours and filtered. The filtrate was evaporated in vacuo and treated with 4N hydrogen chloride in ethyl acetate to afford (2S)-1-[3-hydroxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]amino]-3-phenoxy-2-propanol

hydrochloride (161 mg).

NMR (DMSO- d_6 , δ) : 1.20-1.35 (1H, m), 1.75-2.10 (2H, m), 2.20-2.30 (1H, m), 2.60-2.75 (2H, m), 2.90-3.35 (5H, m), 4.00 (2H, s), 4.22 (1H, br s), 5.90 (1H, d, $J=4.8\text{Hz}$), 6.50-6.70 (2H, m), 6.80-7.05 (4H, m), 7.20-7.45 (2H, m), 9.25 (1H, s)
(+) APCI-MASS (m/z) : 328 (M+H)⁺

Example 147

A mixture of (2S)-1-[N-benzyl-N-(2,3-dimethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol (400 mg) and 10% palladium on activated carbon (50% wet, 100 mg) in methanol (10 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 3 hours and filtered. The filtrate was evaporated in vacuo and treated with 4N hydrogen chloride in ethyl acetate to afford (2S)-1-[2,3-dimethoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-phenoxy-2-propanol hydrochloride (282 mg).

(+) APCI-MASS (m/z) : 372 (M+H)⁺

Example 148

A mixture of (2S)-1-[N-benzyl-N-(1,4-dimethoxy-5,6,8,9-tetrahydro-5H-benzocyclohepten-7-yl)amino]-3-phenoxy-2-propanol (70 mg) and 10% palladium on activated carbon (50% wet, 10 mg) in methanol (10 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 3 hours and filtered. The filtrate was evaporated in vacuo and treated with 4N hydrogen chloride in ethyl acetate to afford (2S)-1-[1,4-dimethoxy-5,6,8,9-tetrahydro-5H-benzocyclohepten-7-yl)amino]-3-phenoxy-2-propanol hydrochloride (446 mg).

(+) APCI-MASS (m/z) : 408 (M+H)⁺

Example 149

A mixture of (2S)-1-[N-benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-5-yl)amino]-3-phenoxy-2-propanol (750 mg) and 10% palladium on activated carbon (50% wet, 200 mg) in methanol (10 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 3 hours and filtered. The filtrate was evaporated in vacuo and treated with 4N hydrogen chloride in ethyl acetate to afford (2S)-1-[3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-5-yl)amino]-3-phenoxy-2-propanol hydrochloride (100 mg).

(+) APCI-MASS (m/z) : 342 (M+H)⁺

Example 150

To a solution of 2-[8-[N-tert-butoxycarbonyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid (100 mg), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (61 mg) and 1-hydroxybenzotriazole (33 mg) in dichloromethane (10 ml) was added 5-aminotetrazole monohydrate (19.3 mg). The resulting mixture was stirred at room temperature for 18 hours and extracted with ethyl acetate. The extract was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel and treated with 4N hydrogen chloride in ethyl acetate to afford 2-[8-[(2S)-2-hydroxy-3-phenoxypropyl)-amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-2-(tetrazol-5-yl)acetamide (50 mg).

NMR (DMSO-d₆, δ) : 1.60-2.10 (3H, m), 2.20-2.30 (1H, m), 2.60-3.60 (7H, m), 3.80-4.30 (2H, m), 4.71 (1H, s), 4.85 (1H, s), 6.60-7.40 (8H, m)

Example 151

To a solution of guanidine hydrochloride (463 mg) in N,N-dimethylformamide (20 ml) was added 28% sodium ethoxide (0.22 ml) under ice-cooling, and the mixture was stirred at

room temperature for 5 minutes. To the resulting mixture was added 2-[8-[(2S)-2-hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid ethyl ester (400 mg). After being stirred at room temperature for 2 hours, the precipitates were collected by filtration and treated with 4N hydrogen chloride in ethyl acetate to afford N-[2-[8-[(2S)-2-hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetyl]guanidine hydrochloride (243 mg).

(+) APCI-MASS (m/z) : 427 (M+H)⁺

Example 152

A mixture of N-[8-[N-benzyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylmethyl]urea (160 mg) and 10% palladium on activated carbon (50% wet, 50 mg) in methanol (10 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 3 hours and filtered. The filtrate was evaporated in vacuo and treated with 4N hydrogen chloride in ethyl acetate to afford N-[8-[(2S)-2-hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylmethyl]urea hydrochloride (100 mg).

(+) APCI-MASS (m/z) : 384 (M+H)⁺

Example 153

A mixture of 1-[8-[N-benzyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylmethyl]-3-butylurea (105 mg) and 10% palladium on activated carbon (50% wet, 50 mg) in methanol (10 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 3 hours and filtered. The filtrate was evaporated in vacuo and treated with 4N hydrogen chloride in ethyl acetate to afford 1-butyl-3-[8-(2S)-2-hydroxy-3-phenoxypropyl]amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylmethyl]urea hydrochloride (80 mg).

(+) APCI-MASS (m/z) : 440 (M+H)⁺

Example 154

A mixture of 3-[[8-[N-benzyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylmethyl]amino]propionic acid tert-butyl ester (100 mg) and 10% palladium on activated carbon (50% wet, 10 mg) in methanol (10 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 3 hours and filtered. The filtrate was evaporated in vacuo and treated with 4N hydrogen chloride in ethyl acetate to afford 3-[[8-[(2S)-2-hydroxy-3-phenoxypropylamino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-ylmethyl]amino]propionic acid dihydrochloride (80 mg).

NMR (DMSO-d₆, δ) : 1.10-1.40 (1H, m), 1.70-2.10 (3H, m), 2.15-3.30 (13H, m), 3.95-4.25 (3H, m), 6.80-7.40 (8H, m), 8.63 (1H, br s)

Example 155

A mixture of 2-[8-[N-benzyl-N-[(2S)-2-hydroxy-3-[4-benzyloxy-3-methanesulfonylamino]phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-butylacetamide (160 mg) and 10% palladium on activated carbon (50% wet, 30 mg) in methanol (10 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 3 hours and filtered. The filtrate was evaporated in vacuo and treated with 4N hydrogen chloride in ethyl acetate to afford N-butyl-2-[8-[[[(2S)-2-hydroxy-3-[4-hydroxy-3-methanesulfonylamino]phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetamide hydrochloride (446 mg).

NMR (DMSO-d₆, δ) : 0.85 (3H, t, J=7.1Hz), 1.20-1.50 (5H, m), 1.60-2.10 (2H, m), 2.22-2.30 (1H, m), 2.70-2.80 (1H, m), 2.94 (3H, s), 3.05-3.30 (7H, m), 3.90 (2H, d, J=4.5Hz), 4.10-4.25 (1H, m), 4.41 (2H,

s), 6.60-6.80 (2H, m), 6.85-6.95 (3H, m), 7.05 (1H, d, J=8.2Hz), 8.04 (1H, br s)

(+) APCI-MASS (m/z) : 480 (M+H)⁺

5 Example 156

A mixture of (2S)-1-[N-benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-3-(4-benzyloxy-3-methanesulfonylamino)phenoxy-2-propanol (750 mg) and 10% palladium on activated carbon (50% wet, 200 mg) in methanol (10 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 3 hours and filtered. The filtrate was evaporated in vacuo and treated with 4N hydrogen chloride in ethyl acetate to afford (2S)-3-[3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl]amino-1-(4-hydroxy-3-methanesulfonylamino)phenoxy-2-propanol hydrochloride (446 mg).

(+) APCI-MASS (m/z) : 342 (M+H)⁺

Example 157

To a solution of 2-[8-[N-tert-butoxycarbonyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid (200 mg), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (122 mg) and 1-hydroxybenzotriazole hydrate (66 mg) in dichloromethane (20 ml) was added dimethylamine hydrochloride (38.4 mg), and the mixture was stirred at room temperature for 18 hours. The resulting mixture was diluted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed (hexane-ethyl acetate) over silica gel and treated with 4N hydrogen chloride in ethyl acetate to afford 2-[8-[[[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N,N-dimethylacetamide (100 mg).

NMR (DMSO-d₆, δ) : 1.10-1.40 (1H, m), 1.60-2.10 (2H,

m), 2.20-2.30 (1H, m), 2.55-3.30 (13H, m), 3.90-4.30 (3H, m), 4.70 (2H, s), 6.89-7.35 (8H, m)
(+) APCI-MASS (m/z) : 413 (M+H)⁺

5 Example 158 to Example 201

Coupling of 2-[8-[N-tert-butoxycarbonyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid with n different type of amines

10.


To a solution of 2-[8-[N-tert-butoxycarbonyl-N-[(2S)-2-hydroxy-3-phenoxypropyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid (Starting compound) (n x 0.01 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (n x 0.015 mmol) and 1-hydroxybenzotriazole (n x 0.012 mmol) in N,N'-dimethylformamide (n x 0.1 ml) was added 1M N,N-dimethylformamide solution of an amine (0.012 mmol) [n different type of amines] and stirred at room temperature for 2 hours.

20


To each reaction mixture was added 5% sodium hydrogencarbonate solution (0.40 ml), following by extraction with ethyl acetate (0.35 ml). The resultant aqueous layer was further extracted with ethyl acetate (0.2 ml). The combined organic layer was washed with water (0.30 ml). Then the resultant aqueous layer was additionally extracted with ethyl acetate (0.2 ml x 2). The combined organic layer was concentrated by nitrogen flow and treated with 4N hydrogen chloride in ethyl acetate and concentrated by nitrogen flow. The resultant residue was dissolved in dimethylsulfoxide (1.0 ml) to give a ca. 1.0×10^{-2} M dimethylsulfoxide solution of the object compound which was subjected to analysis by mass spectrum.

30

35














(Starting compound)



















(Object compound)











The formula $-N \begin{smallmatrix} R \\ R' \end{smallmatrix}$ of the Object compounds and their mass spectrums in the following Examples 158 to Example 201 are shown in Table 1.







| Example No. | $-N \begin{smallmatrix} R \\ R' \end{smallmatrix}$ | MALDI-MS (m/z) |
|-------------|--|------------------------|
| 158 | | 457 (M+H) ⁺ |
| 159 | | 455 (M+H) ⁺ |
| 160 | | 443 (M+H) ⁺ |
| 161 | | 485 (M+H) ⁺ |

| Example No. |  | MALDI-MS (m/z) |
|-------------|---|------------------------|
| 162 |  | 443 (M+H) ⁺ |
| 163 |  | 498 (M+H) ⁺ |
| 164 |  | 467 (M+H) ⁺ |
| 165 |  | 543 (M+H) ⁺ |
| 166 |  | 498 (M+H) ⁺ |
| 167 |  | 490 (M+H) ⁺ |
| 168 |  | 476 (M+H) ⁺ |
| 169 |  | 493 (M+H) ⁺ |
| 170 |  | 517 (M+H) ⁺ |
| 171 |  | 558 (M+H) ⁺ |

| Example No. | $\begin{array}{c} \text{R} \\ \\ -\text{N}- \\ \\ \text{R}' \end{array}$ | MALDI-MS (m/z) |
|-------------|---|------------------------|
| 172 |  | 573 (M+H) ⁺ |
| 173 |  | 517 (M+H) ⁺ |
| 174 |  | 529 (M+H) ⁺ |
| 175 |  | 506 (M+H) ⁺ |
| 176 |  | 479 (M+H) ⁺ |
| 177 |  | 532 (M+H) ⁺ |
| 178 |  | 620 (M+H) ⁺ |
| 179 |  | 565 (M+H) ⁺ |

| Example No. |  | MALDI-MS (m/z) |
|-------------|---|------------------------|
| 180 |  | 680 (M+H) ⁺ |
| 181 |  | 623 (M+H) ⁺ |
| 182 |  | 489 (M+H) ⁺ |
| 183 |  | 565 (M+H) ⁺ |
| 184 |  | 542 (M+H) ⁺ |
| 185 |  | 504 (M+H) ⁺ |
| 186 |  | 475 (M+H) ⁺ |

| Example No. |  | MALDI-MS (m/z) |
|-------------|---|------------------------|
| 187 |  | 491 (M+H) ⁺ |
| 188 |  | 495 (M+H) ⁺ |
| 189 |  | 468 (M+H) ⁺ |
| 190 |  | 487 (M+H) ⁺ |
| 191 |  | 527 (M+H) ⁺ |
| 192 |  | 455 (M+H) ⁺ |
| 193 |  | 455 (M+H) ⁺ |
| 194 |  | 469 (M+H) ⁺ |
| 195 |  | 483 (M+H) ⁺ |

| Example No. | $\begin{array}{c} \text{R} \\ \\ -\text{N}- \\ \\ \text{R}' \end{array}$ | MALDI-MS (m/z) |
|-------------|---|------------------------|
| 196 |  | 479 (M+H) ⁺ |
| 197 |  | 504 (M+H) ⁺ |
| 198 |  | 479 (M+H) ⁺ |
| 199 |  | 551 (M+H) ⁺ |
| 200 |  | 503 (M+H) ⁺ |
| 201 |  | 481 (M+H) ⁺ |

Example 202

Under nitrogen, a solution of 2-(8-amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy)-N,N-dimethylacetamide (129 mg), N-[2-benzyloxy-5-[(1R)-2-iodo-1-(triethylsilyloxy)-

ethyl]phenyl]methanesulfonamide (280 mg) and N,N-diisopropylethylamine (0.34 ml) in 1,3-dimethyl-2-imidazolidinone (5 ml) was stirred at 110°C for 42 hours. The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous sodium sulfate and evaporated in vacuo. To the residue in ethyl acetate (4 ml) was added 4N hydrogen chloride in ethyl acetate (1 ml), and the mixture was stirred at room temperature for 1.5 hours. After evaporation, the residue was dissolved into a mixture of saturated aqueous sodium hydrogencarbonate and ethyl acetate, followed by separation. The organic layer was washed with brine, dried over anhydrous magnesium sulfate, and evaporated in vacuo. The residue was purified by column chromatography on silica gel (chloroform:methanol = 50:1 to 10:1) to give 2-[8-[(2R)-2-(4-benzyloxy-3-methanesulfonylamino)phenyl-2-hydroxyethylamino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N,N-dimethylacetamide (145 mg).

NMR (CHCl₃, δ) : 1.4-2.1 (4H, m), 2.55-3.15 (7H, m),
 2.91 (3H, s), 2.96 (3H, m), 3.08 (3H, m), 4.45-4.7
 (3H, m), 5.10 (2H, s), 6.65-6.8 (2H, m), 6.9-7.0
 (2H, m), 7.1-7.2 (1H, m), 7.35-7.55 (6H, m)
 (+) APCI-MASS (m/z) : 582 (M+H)⁺

The following compound was obtained according to a similar manner to that of Example 202.

Example 203

2-[8-[(2R)-2-Hydroxy-2-(4-hydroxy-3-methanesulfonylaminophenyl)ethylamino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N,N-dimethylacetamide hydrochloride

NMR (DMSO-d₆, δ) : 1.1-1.4 (1H, m), 1.7-2.05 (2H, m),
 2.2-2.4 (1H, m), 2.55-3.2 (7H, m), 2.83 (3H, s),
 2.95 (3H, s), 2.99 (3H, s), 4.73 (2H, s), 4.8-4.95

(1H, m), 6.67 (1H, dd, J=2.2 and 8.2Hz), 6.75-6.9
 (1H, m), 6.9-7.4 (4H, m)
 (+) APCI-MASS (m/z) : 492 (M-HCl+H)⁺

Example 204

Under nitrogen, to a solution of N-[3-[(1S)-2-amino-1-hydroxyethyl]-4-fluorophenyl]methanesulfonamide (200 mg) and 3-methoxy-6,7,8,9-tetrahydrobenzocyclohepten-6-one (146 mg) in tetrahydrofuran (10 ml) was added sodium triacetoxymethylborohydride (257 mg) at room temperature, and the mixture was stirred at the same temperature for 24 hours. The resulting mixture was poured into saturated aqueous sodium bicarbonate solution and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and evaporated in vacuo. The residue was treated with 4N hydrogen chloride in ethyl acetate to afford N-[4-fluoro-3-[(1S)-1-hydroxy-2-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-ylamino)ethyl]phenyl]-methanesulfonamide hydrochloride (304 mg).

(+) APCI-MASS (m/z) : 423 (M+H)⁺

Example 205

Under nitrogen, a solution of 3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-ylamine (720 mg), N-[2-benzyloxy-5-[(1R)-2-iodo-1-(triethylsilyloxy)ethyl]phenyl]-methanesulfonamide (2.1 g) and N,N-diisopropylethylamine (2.6 ml) in N,N-dimethylacetamide (10 ml) was stirred at 110°C for 96 hours. The resulting mixture was poured into aqueous 10% sodium hydrogensulfite and extracted with ethyl acetate. The organic layer was successively washed with saturated aqueous sodium hydrogencarbonate and brine, dried over anhydrous sodium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (chloroform:methanol = 40:1 to 20:1) to give N-[2-benzyloxy-5-[(1R)-1-hydroxy-2-(3-methoxy-6,7,8,9-tetrahydro-5H-benzo-

cyclohepten-6-ylamino)ethyl]phenyl]methanesulfonamide (381 mg).

NMR (CDCl_3 , δ) : 1.4-2.1 (5H, m), 2.5-3.1 (7H, m),
 2.90 (3H, s), 3.77 (3H, s), 4.5-4.6 (1H, m), 5.10
 (2H, s), 6.6-6.75 (2H, m), 6.97 (1H, d, $J=8.5\text{Hz}$),
 6.99 (1H, d, $J=8.0\text{Hz}$), 7.1-7.2 (1H, m), 7.3-7.5
 (5H, m), 7.50 (1H, d, $J=1.8\text{Hz}$)
 (+) APCI-MASS (m/z) : 511 (M+H)⁺

The following compound was obtained according to a similar manner to that of Example 132.

Example 206

5 N-[2-Hydroxy-5-[(1R)-1-hydroxy-2-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-ylamino)ethyl]phenyl]-methanesulfonamide hydrochloride

10 NMR ($\text{DMSO}-d_6$, δ) : 1.05-1.4 (1H, m), 1.7-2.1 (2H, m),
 2.2-2.4 (1H, m), 2.6-2.8 (2H, m), 2.9-3.3 (5H, m),
 2.95 (3H, s), 3.71 (3H, s), 4.8-4.95 (1H, m), 6.65-
 7.2 (5H, m), 7.28 (1H, s)
 (+) APCI-MASS (m/z) : 421 (M-HCl+H)⁺

Example 207

15 Under nitrogen, a solution of 2-(8-amino-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy)acetic acid tert-butyl ester (321 mg), N-[2-benzyloxy-5-[(1R)-2-iodo-1-(triethylsilyloxy)ethyl]phenyl]methanesulfonamide (620 mg) and N,N-diisopropylethylamine (0.77 ml) in 1,3-dimethyl-2-imidazolidinone (5 ml) was stirred at 110°C for 24 hours.
 20 The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate and extracted with ethyl acetate. The organic layer was successively washed with water twice and brine, dried over anhydrous sodium sulfate and evaporated
 25 in vacuo. Under nitrogen, to the residue in tetrahydrofuran (10 ml) was added 1M tetra-n-butylammonium fluoride in

tetrahydrofuran (2 ml) at 5°C, and the mixture was stirred at room temperature for 1 hour. The resulting mixture was poured into saturated aqueous sodium hydrogencarbonate and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (chloroform:methanol = 50:1) to give 2-[8-[(2R)-2-(4-benzyloxy-3-methanesulfonylamino)phenyl]-2-hydroxyethylamino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid tert-butyl ester (366 mg).

(+) APCI-MASS (m/z) : 611 (M+H)⁺

Example 208

A mixture of 2-[8-[(2R)-2-(4-benzyloxy-3-methanesulfonylamino)phenyl]-2-hydroxyethylamino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid tert-butyl ester (353 mg) and 10% palladium on activated carbon (50% wet, 100 mg) in methanol (5 ml) was stirred at room temperature in the presence of hydrogen at an atmospheric pressure for 3.5 hours. After filtration, the filtrate was evaporated in vacuo. The residue was purified by column chromatography on silica gel (chloroform:methanol = 10:1 to 5:1) to give 2-[8-[(2R)-2-hydroxy-2-(4-hydroxy-3-methanesulfonylamino)phenyl]-ethylamino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid tert-butyl ester (268 mg).

(+) APCI-MASS (m/z) : 521 (M+H)⁺

Example 209

Under nitrogen, to a solution of 2-[8-[(2R)-2-hydroxy-2-(4-hydroxy-3-methanesulfonylamino)phenyl]ethylamino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]acetic acid tert-butyl ester (257 mg) in dichloromethane (5 ml) was added trifluoroacetic acid (1 ml) at 5°C, and the mixture was stirred at room temperature for 3 hours. The resulting mixture was evaporated in vacuo. The residue was treated

with 4N hydrogen chloride in 1,4-dioxane to give 2-[8-[(2R)-2-hydroxy-2-(4-hydroxy-3-methanesulfonylamino)phenyl)-ethylamino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-acetic acid hydrochloride (247 mg).

5 (+) APCI-MASS (m/z) : 465 (M-HCl+H)⁺

The following compound was obtained according to a similar manner to that of Example 205.

10 Example 210

2-[8-[(2R)-2-(4-Benzyloxy-3-methanesulfonylamino)phenyl)-2-hydroxyethylamino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-phenylacetamide

15 NMR (CDCl₃, δ) : 1.4-2.1 (4H, m), 2.5-3.1 (7H, m), 2.90 (3H, s), 4.5-4.65 (1H, m), 4.58 (2H, s), 5.09 (2H, s), 6.71 (1H, dd, J=2.6 and 8.1Hz), 6.81 (1H, d, J=2.6Hz), 6.9-7.2 (4H, m), 7.25-7.65 (10H, m), 8.32 (1H, br s)

20 (+) APCI-MASS (m/z) : 630 (M+H)⁺

The following compound was obtained according to a similar manner to that of Example 79.

25 Example 211

2-[8-[(2R)-2-Hydroxy-2-(4-hydroxy-3-methanesulfonylamino)phenyl)ethylamino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-phenylacetamide hydrochloride

30 NMR (DMSO-d₆, δ) : 1.1 (4H, m), 2.6-2.8 (2H, m), 2.9-3.7 (8H, m), 4.67 (2H, s), 4.8-4.9 (1H, m), 6.7-7.4 (9H, m), 7.65 (2H, d, J=7.8Hz)

(+) APCI-MASS (m/z) : 540 (M-HCl+H)⁺

35 Example 212

Under nitrogen, a solution of 3-oxiranylpiperidine (1.9 g) and N-benzyl-(3-methoxy-6,7,8,9-tetrahydro-5H-

benzocyclohepten-6-yl)amine (3.3 g) in ethanol (15 ml) was refluxed for 5 hours. After removal of ethanol in vacuo, the residue was dissolved into a mixture of aqueous sodium hydrogencarbonate and ethyl acetate. After separation, the organic layer was washed with brine, dried over anhydrous magnesium sulfate and evaporated in vacuo. The residue was purified by column chromatography on silica gel (chloroform:methanol = 20:1) to give 2-[N-benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-1-(pyridin-3-yl)ethanol (520 mg).

(+) APCI-MASS (m/z) : 403 (M+H)⁺

The following compound was obtained according to a similar manner to that of Example 132.

Example 213

2-(3-Methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino-1-(pyridin-3-yl)ethanol dihydrochloride

(+) APCI-MASS (m/z) : 313 (M-2HCl+H)⁺

Example 214

A mixture of N-benzyl-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amine (0.30 g) and (R)-3-chlorophenyl oxirane (0.18 g) in ethanol (0.6 ml) was refluxed overnight. The reaction mixture was evaporated in vacuo, and the residue was purified by column chromatography on silica gel eluting with a mixture of hexane and ethyl acetate (5:1). The product (0.43 g) was treated with 4N hydrogen chloride in ethyl acetate and powdered from diisopropyl ether to afford (1R)-2-[N-benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-1-(3-chlorophenyl)ethanol (0.40 g).

NMR (DMSO-d₆, δ) : 1.10-1.25 (1H, m), 1.91-2.18 (2H, m), 2.42-2.80 (3H, m), 2.98-3.62 (5H, m), 3.74 (3H, s), 4.43-4.73 (3H, m), 6.36-6.46 (1H, br), 6.68

(1H, d, J=7.9Hz), 7.01 (1H, d, J=8.3Hz), 7.11-7.19
(1H, m), 7.29-7.55 (7H, m), 7.68-7.90 (2H, m),
10.00-10.15 (1H, br)

(+) APCI-MASS (m/z) : 436, 438 (M+H)⁺

5

Example 215

A mixture of (1R)-2-[N-benzyl-N-(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)amino]-1-(3-chlorophenyl)ethanol hydrochloride (0.20 g), chlorobenzene
10 (0.86 ml) and 10% palladium on carbon (0.01 g) in methanol (4 ml) was stirred at room temperature under hydrogen atmosphere for 1.5 hours. The catalyst was filtered off and washed with methanol. The filtrate was evaporated in vacuo, and the residue was powdered from diisopropyl ether to afford (1R)-2-
15 [(3-methoxy-6,7,8,9-tetrahydro-5H-benzocyclohepten-6-yl)-amino]-1-(3-chlorophenyl)ethanol hydrochloride (0.14 g).

IR (KBr) : 3328, 2933, 1608, 1502, 1444, 1261 cm⁻¹

NMR (DMSO-d₆, δ) : 1.17-1.35 (1H, m), 1.78-2.08 (2H, m), 2.22-2.37 (1H, m), 2.62-2.72 (2H, m), 2.98-3.30
20 (5H, m), 3.71 (3H, s), 4.95-5.05 (1H, m), 6.33 (1H, d, J=4.0Hz), 6.70 (1H, d, J=8.2Hz), 6.83 (1H, dd, J=12.3, 2.5Hz), 7.04 (1H, d, J=8.2Hz), 7.30-7.43 (3H, m), 7.52 (1H, s), 8.72 (1H, br s), 9.00 (1H, br s)

25

(+) APCI-MASS (m/z) : 346, 348 (M-HCl+H)⁺

The following compound was obtained according to a similar manner to that of Example 214.

30

Example 216

2-[8-[N-Benzyl-N-[(2R)-2-(3-chlorophenyl)-2-hydroxyethyl]amino]-6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-phenylacetamide hydrochloride

NMR (DMSO-d₆, δ) : 1.05-1.30 (1H, m), 1.91-2.15 (2H, m), 2.40-2.50 (1H, m), 2.62-2.75 (2H, m), 3.02-3.60
35

(6H, m), 4.45-4.72 (4H, m), 6.34-6.45 (1H, br),
6.75 (1H, d, J=8.2Hz), 7.01-7.20 (3H, m), 7.27-7.48
(9H, m), 7.62-7.68 (2H, m), 7.70-7.90 (2H, m), 9.83
(1H, br s), 10.19 (1H, d, J=7.8Hz)

5 (+) APCI-MASS (m/z) : 555, 557 (M+H)⁺

The following compound was obtained according to a
similar manner to that of Example 215.

10 Example 217

2-[8-[[(2R)-2-(3-Chlorophenyl)-2-hydroxyethyl]amino]-
6,7,8,9-tetrahydro-5H-benzocyclohepten-2-yloxy]-N-
phenylacetamide

15 IR (KBr) : 3394, 2933, 1679, 1600, 1540, 1502, 1442,
1253 cm⁻¹

20 NMR (DMSO-d₆, δ) : 1.16-1.30 (1H, m), 1.72-2.05 (2H,
m), 2.22-2.38 (1H, m), 2.60-2.70 (2H, m), 3.00-3.35
(5H, m), 4.66 (2H, d, J=2.4Hz), 4.95-5.04 (1H, m),
6.33 (1H, br s), 6.77 (1H, d, J=8.4Hz), 6.92 (1H,
25 d, J=12.3Hz), 7.04-7.10 (2H, m), 7.27-7.41 (5H, m),
7.52 (1H, s), 7.62-7.66 (2H, m), 8.75 (1H, br s),
9.12 (1H, br s), 10.16 (1H, d, J=6.1Hz)

(+) ESI-MASS (m/z) : 465, 467 (M+H)⁺

25 Example 218

According to a similar manner to that of Test 1
described before, the effects of the present compounds on the
intravesical pressure were evaluated. Particularly,
ED50 (μg/kg) of the compound prepared in Example 155 was 10.8.